# Annual Deer Population Report & 2021-22 Antlerless License Allocations Recommendations



March 31, 2021

Pennsylvania Game Commission Bureau of Wildlife Management Deer and Elk Section

# Summary of 2021-22 Antlerless Allocations to Achieve Deer Plan Goals

WMU	Population Trend	Deer Plan Population Objective	2020-21 Approved Allocation	2021-22 Deer Plan Recommendation (14-day Concurrent)	Comments
1A	Stable	Stabilize	49,000	40,000	Change of 1 week to 2 weeks concurrent.
1B	Stable	Stabilize	41,000	32,000	Change of 1 week to 2 weeks concurrent.
2A	Stable	Stabilize	46,000	39,000	Change of 1 week to 2 weeks concurrent.
2B	Stable	Stabilize	49,000	49,000	No change.
2C	Stable	Reduce	58,000	67,000	Maintaining the increased harvest by 1.5 antlerless deer/mi <sup>2</sup> to reduce population because of CWD.
2D	Stable	Reduce	60,000	74,000	Maintaining the increased harvest by 1.5 antlerless deer/mi <sup>2</sup> to reduce population because of CWD.
2E	Stable	Reduce	39,000	42,000	Maintaining the increased harvest by 1.5 antlerless deer/mi <sup>2</sup> to reduce population because of CWD.
2F	Increasing	Stabilize	36,000	32,000	Increase harvest by 1.0 antlerless deer/mi <sup>2</sup> to stop population growth. Change of 1 week to 2 weeks concurrent.
2G	Stable	Stabilize	27,000	23,000	Change of 1 week to 2 weeks concurrent.
2H	Increasing	Stabilize	7,000	9,000	Increase harvest by 1.0 antlerless deer/mi <sup>2</sup> to stop population growth. Change of 1 week to 2 weeks concurrent.
3A	Stable	Stabilize	21,000	19,000	Change of 1 week to 2 weeks concurrent.
3B	Stable	Stabilize	33,000	30,000	Change of 1 week to 2 weeks concurrent.
3C	Stable	Stabilize	49,000	33,000	Change of 1 week to 2 weeks concurrent.
3D	Increasing	Reduce	36,000	36,000	Increase harvest by 1.0 antlerless deer/mi <sup>2</sup> to reduce population because of forest impacts. Change of 1 week to 2 weeks concurrent.
4A	Stable	Reduce	49,000	50,000	Increase harvest by 2.0 antlerless deer/mi <sup>2</sup> to reduce population because of CWD (CWD prevalence goals are not being met in the established area).
4B	Decreasing	Reduce	33,000	34,000	Maintaining the increased harvest by 1.5 antlerless deer/mi <sup>2</sup> to reduce population because of CWD.
4C	Stable	Stabilize	32,000	29,000	Change of 1 week to 2 weeks concurrent.
4D	Stable	Reduce	45,000	55,000	Maintaining the increased harvest by 1.5 antlerless deer/mi <sup>2</sup> to reduce population because of CWD.
4E	Stable	Reduce	37,000	42,000	Increase harvest by 1.5 antlerless deer/mi <sup>2</sup> to reduce population because of forest impacts and CWD (new). Change of 1 week to 2 weeks concurrent.
5A	Stable	Reduce	26,000	31,000	Maintaining the increased harvest by 1.5 antlerless deer/mi <sup>2</sup> to reduce population because of CWD.
5B	Stable	Stabilize	60,000	60,000	Change of 1 week to 2 weeks concurrent (based on history of this WMU, minimal increase in license efficiency expected).
5C	Stable	Stabilize	70,000	70,000	No change.
5D	Stable	Stabilize	29,000	29,000	No change.

Data presented in this report represent collaborative efforts between the U.S. Forest Service, Pennsylvania's Department of Conservation and Natural Resources, the Pennsylvania Cooperative Fish and Wildlife Research Unit at Penn State University, Responsive Management, and the Game Commission's bureaus of Information and Education, Wildlife Habitat Management, and Wildlife Management. For more information on the deer management program and data and methods used to assess progress towards management goals, visit the Game Commission's website, <a href="www.pgc.pa.gov">www.pgc.pa.gov</a>, to find the "2009-2018 White-tailed Deer Management Plan".

#### **Deer Management Goals**

Deer management goals direct Game Commission staff in formulating deer management recommendations. Current management goals that directly affect antlerless allocations are to manage deer for healthy deer, healthy forest habitat, and acceptable levels of deer-human conflicts. These goals were identified by a group of public stakeholders in 2002 and continue to be supported by a clear majority of Pennsylvania citizens and hunters (Figure 1).




Figure 1. Percent of respondents that agree with deer management goals. The citizen survey was completed by Responsive Management in the fall of 2011, and the deer hunter survey was completed by the Deer and Elk Section and Bureau of Wildlife Management in the fall of 2020.

#### **Step-by-Step Deer Management Recommendation Guide**

The deer management program considers data for each goal to arrive at a deer population recommendation in a defined process (see pages 7 and 8). This process has been revised as new data are incorporated into the program and will continue to evolve as more data and understanding are gained. Decision points (i.e., fawn to doe ratio declining?) are based on published protocols from the wildlife and forestry professions.

#### Do PA residents want fewer or more deer?

This question is answered using results of the survey conducted by Responsive Management of Pennsylvania residents in 2019. If most surveyed residents in a WMU want less deer, the recommendation would be to reduce the deer population. If the deer health goal is met, forest habitat is good, and WMU residents want more deer, the recommendation would be to increase the deer population.

#### Is CWD present in free-ranging deer?

This question is answered using results from the thousands of deer tested annually for chronic wasting disease (CWD). If CWD is present in free-ranging deer, then management recommendations are to stabilize or reduce WMU populations. Additional antlerless deer can be removed using Deer Management Assistance Program permits in accordance with the CWD response plan. Increasing the antlerless harvest serves 2 purposes that are important to efforts to contain CWD; (1) increased antlerless harvest removes more deer from the population and allows the Game Commission to test more deer in our efforts to obtain the best information on the extent of the disease, and (2) increased antlerless harvest can reduce deer populations and spread of CWD.

#### Is fawn to doe ratio declining?

This question is answered using results from the age structure of the antlerless harvest. These data are collected each year by trained Game Commission deer agers from across the state. If the proportion of fawns in the antlerless harvest (hereafter referred to as fawn to doe ratio) is declining and the population is not achieving its objective (i.e., population is declining and objective is to maintain a stable deer population), then the antlerless allocation would be reduced to stop the population decline. The antlerless harvest will have the greatest influence on the population because hunting accounts for most deer mortalities in Pennsylvania. If the fawn to doe ratio is stable or if the population is meeting its objective (i.e., population is stable and objective is stable), no management action is taken.

#### Has deer population been stable or increasing for 6 years?

This question is answered using results from the Pennsylvania Sex-Age-Kill deer population model and deer harvest indices (i.e., antlered harvest, antlerless catch-per-unit-effort). The 6-year time period is necessary because of the 5-year time period to collect the forest data. The sixth

year is added because only  $2^{nd}$  year seedlings are counted in the forest data. As a result, a complete forest data set includes effects of deer from the previous 6 years.

If the deer population is decreasing the recommendation is to stabilize the population at the lower level to see if forest habitat improves given the lower deer population. If the deer population is stable or increasing, the process continues to the next step.

#### Is forest habitat good?

This question is answered using results from the Pennsylvania Regeneration Study. If 70% of forested plots have adequate regeneration, forest habitat is considered good. If less than 50% of forested plots have adequate regeneration, forest habitat is considered poor. If 50% to 70% of forested plots have adequate regeneration, forest habitat is considered fair.

#### Is plot to plot regeneration improving?

This question is answered using results from the Pennsylvania Regeneration Study. In this step, results from individual plots are compared in a paired analysis. For example, plot measurements from 2005 are compared to their remeasured results in 2010 to see if regeneration has improved on individual plots. All plots with 2 measures are included in this analysis. If regeneration is improving, then the deer population trend can be stabilized. If regeneration is not improving, the process continues to the next step.

#### Is plot to plot deer impact improving?

This question is answered using results from the Pennsylvania Regeneration Study. In this step, results from assessments of deer impact on a scale from 1 (very low) to 5 (very high) are compared in the same way as the plot to plot regeneration analysis. If deer impact is improving (i.e., going from a 4 [high] to 3 [moderate]) on enough plots, then the deer population trend can be stabilized. If deer impact is not improving, the process continues to the next step.


#### Is mean deer impact 3 or less?

This question is answered from the Pennsylvania Regeneration Study. In this step, the mean deer impact for all plots measured in the most recent 5-year period is statistically compared to an objective of 3 (i.e., moderate impact). If deer impact is significantly greater than 3 (moderate), then the deer impact is too high and the deer population should be reduced. If deer impact is less than or not different from 3 (moderate) then the deer population trend can be stabilized.

Guides on pages 7 and 8 are used to develop deer population recommendations based on goals and objectives of deer management plan. Recommendation guide for WMUs 2B, 5C, and 5D differs because of lack of forest data in these highly developed WMUs.




# **Deer Management Recommendation Process**





# **Deer Management Recommendation Guide**

FOR WMUs 2B, 5C, and 5D



#### **Step-by-Step Antlerless License Allocation Calculations**

Antlerless allocations are calculated by referring to results from previous seasons. For example, if a population has remained stable with an annual harvest of 3,000 antlerless deer, the same level of harvest would be expected to maintain the stable population. If it has taken 3 antlerless licenses to harvest 1 antlerless deer over the last 3 years, the allocation to stabilize this population would be 3,000 antlerless deer harvested x 3 licenses/antlerless deer harvested = 9,000 antlerless licenses.


Bold numbers in Table 1 below were the ones used in the calculations for 2021-22 allocations. For WMUs 2C, 2D, 2E, 4A, 4B, 4D, and 5A, the concurrent season was increased from 1 week to 2 weeks in 2020, thus estimates of licenses/deer from the 2020 season were used instead of the 3 year mean for 2021 allocation calculations.

Table 1. Antlerless licenses needed to harvest 1 antlerless deer (license/deer) based on historic results for each WMU, March 2021.

WMU	2018-19	2019-20	2020-21	3-year Average
1A	3.8	3.7	2.8	3.4
1B	2.4	2.8	2.3	2.5
2A	4.2	4.4	3.9	4.1
2B	3.8	4.3	3.3	3.8
2C	4.0	4.1	3.7	3.9
2D	3.1	3.8	3.2	3.4
2E	3.1	4.2	3.4	3.6
2F	3.1	3.5	3.6	3.4
2G	4.1	4.3	4.0	4.1
2H	3.3	5.6	4.5	4.5
3A	3.0	3.5	3.1	3.2
3B	3.5	3.7	3.9	3.7
3C	3.1	3.6	3.4	3.4
3D	4.4	5.1	5.7	5.1
4A	5.5	5.8	4.0	5.1
4B	3.9	4.4	3.1	3.8
4C	4.2	4.3	4.0	4.2
4D	3.9	4.5	3.7	4.0
4E	3.4	3.6	3.3	3.4
5A	4.9	4.4	4.3	4.5
5B	4.1	4.5	3.6	4.1
5C	4.2	4.8	4.6	4.6
5D	4.6	4.3	4.4	4.4

### Trend in Fawn to Doe Ratios, 2015 to 2020

(Supporting data in WMU worksheets, pages 24 to 69)



#### Legend


Decreasing Fawn to Doe Ratio Stable Fawn to Doe Ratio Increasing Fawn to Doe Ratio





# Forest Regeneration, 2015 to 2019

(Supporting data in WMU worksheets, pages 24 to 69)



#### Legend


Poor Forest Regeneration Levels Fair Forest Regeneration Levels Good Forest Regeneration Levels





# Plot to Plot Change in Regeneration, 5-year Change

(Supporting data in WMU worksheets, pages 24 to 69)



#### Legend


Declining Regeneration
No Change in Regeneration
Improving Regeneration





# Deer Impact Level, 2015 to 2019

(Supporting data in WMU worksheets, pages 24 to 69)



#### Legend

Deer Impact is Too High (> 3)


Deer Impact is Acceptable (3 or less)





# Plot to Plot Change in Deer Impact, 5-year Change

(Supporting data in WMU worksheets, pages 24 to 69)



#### Legend


Increasing Deer Impact
No Change in Deer Impact
Improving Deer Impact





# Post-Hunt Deer Population Trends, 2016 to 2021

(Supporting data in WMU worksheets, pages 24 to 69)



#### Legend


Declining Deer Population Stable Deer Population Increasing Deer Population





# Pennsylvania Residents Opinions on Deer Populations, 2019

(Supporting data in WMU worksheets, pages 24 to 69)



#### Legend


Most Residents Say Deer Population Too High Most Residents Say Deer Population Just Right Most Residents Say Deer Population Too Low









# Residents Opinions on Deer Populations 2011 vs. 2019






#### Legend


More than 25% say Deer Population Too High Less than 25% say Deer Population Too High and less than 25% say Too Low More than 25% say Deer Population Too Low





# Deer Hunters Opinions on Deer Populations 2011 vs. 2020





#### Legend


More than 25% say Deer Population Too High Less than 25% say Deer Population Too High and less than 25% say Too Low More than 25% say Deer Population Too Low







# **Chronic Wasting Disease (CWD), March 2021**



#### Legend


WMUs with CWD Detected in Wild Deer WMUs with No CWD Positive Wild Deer Detected





#### 2020-21 Regular Firearms Seasons

In 2001, a 12-day concurrent antlered and antlerless firearms season began. The objectives of this longer antlerless season were to give hunters more time to hunt antlerless deer and to create a more consistent harvest from year to year. Antlerless allocations in each WMU determined antlerless harvest. Beginning in 2008, some WMUs were changed to a 5-day antlered only season followed by a 7-day concurrent antlered and antlerless season. In 2010, 2011, 2014, 2015, and 2017 additional WMUs were changed to the 5/7 season format. By 2019, only WMUs 2B, 5C, and 5D had a two-week concurrent antlered and antlerless firearms season.



#### Legend

7-day Antlered Only and 7-day Antlered and Antlerless Concurrent Firearms Season 14-day Antlered and Antlerless Concurrent Firearms Season








# **2020-21 Antlered Deer Harvest Density**

(Antlered deer harvested per square mile of area)



#### Legend

Less than 2 antlered deer per square mile

2.0 to 3.0 antlered deer per square mile

3.1 to 4.0 antlered deer per square mile

More than 4.0 antlered deer per square mile






# Recommendation Guides and Deer Population Datasheets

Recommendation guides (see pages 7 and 8) provide a step-by-step progression through the deer plan goals and measurable objectives to arrive at a deer population recommendation.

Supporting data for these guides are found in the individual WMU datasheets that follow.

#### **WMU Antlerless Allocation Worksheets**

#### **Example**



#### 2021-22 Pennsylvania Game Commission Antlerless Allocation Worksheet

 WMU
 1A
 % Developed
 % Forest
 %Ag/Field
 %Public
 Area (sq mi)

 9%
 45%
 36%
 3%
 1,846

er Harve	est													
							Antle	red Ha	rvest					
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12.000									_
2005	5,468	13,427	40,000	2.9	12,000 -									
2006	5,791	13,214	42,000	3.1										
2007	4,896	12,490	42,000	3.3	10,000 -									
2008	5,392	12,611	42,000	3.3										ı
2009	5,500	10,689	42,000	4.0	8,000 -									-
2010	5,918	11,940	41,705	3.5										ı
2011	5,171	9,839	42,000	4.3	6,000 -			_	_					Ш
2012	6,078	11,859	42,000	3.5	0,000			l _		_				
2013	6,420	13,915	49,000	3.5										
2014	5,131	10,792	47,000	4.4	4,000 -									
2015	6,031	9,122	46,000	5.0										
2016	6,500	10,377	46,000	4.4	2,000 -									$\vdash$
2017	6,279	12,612	52,000	4.1										
2018	5,802	12,442	48,000	3.8		<u> </u>				, <b>.</b> ,		<b>,</b> ■,		Ļ
2019	6,416	13,160	49,000	3.7		st top toy tog	108 2015	,202,20	y 3	2h 3	\$ %	2027 20	3 203	£
2020	9,210	17,509	49,000	2.8	γ γ	かかかく	\$ \$0.	\$ \$	20.	\$ \$	, 5.	\$ \$	20.	₯.

RED=7 day season


POST-HUNT	Deer Population		Trend	Stable
Year	Total	[		
2008	34,007	160,000		
2009	36,152	140,000		
2010	44,148	1.0,000		
2011	41,549	120,000		
2012	42,420			
2013	48,472	100,000		
2014	55,114	80,000		
2015	49,169	80,000		
2016	62,237	60,000		
2017	65,707			
2018	53,244	40,000		
2019	46,208	20.000		
2020	51,804	20,000		
2021	99,568	0	<u> </u>	<b>■</b> , <b>■</b> , <b>■</b> , <b>■</b> ,
		" " " " " " " " " " " " " " " " " " "	on son son son s	93 203 200 2012

Declining

#### DEER HEALTH: Fawn to Doe Ratio<sup>2</sup>

**Trend** 

Year	Total
2008	0.44
2009	0.45
2010	0.46
2011	0.43
2012	0.48
2013	0.41
2014	0.46
2015	0.42
2016	0.43
2017	0.40
2018	0.39
2019	0.37
2020	0.35




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

**Regeneration Assessment** Good

Year	% Adequate
2003-07	53%
2004-08	54%
2005-09	55%
2006-10	57%
2007-11	55%
2008-12	53%
2009-13	57%
2010-14	55%
2011-15	54%
2012-16	53%
2013-17	50%
2014-18	64%
2015-19	67%



Citizen Survey	/ Results	2019	(2011)	)

Too High 26%(16%) Just Right 55%(54%) **Too Low** 13%(23%)

Antlerless Allocation Recommendation								
	Increase	Stable	Decrease					
Firearm Season Option	Harvest	Harvest	Harvest					
7 day antlered & 7 day concurrent	56,000	50,000	43,000					
14 day concurrent	45,000	40,000	35,000					

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.

#### 2021-22 Pennsylvania Game Commission Antlerless Allocation Worksheet


			VVIV	io Characteri	Stics	
WMU	1B	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		7%	54%	32%	4%	2.115

								Antl	ered F	larves	t			
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	13.000									
2005	6,382	10,707	27,000	2.5	12,000 -									
2006	6,773	11,974	30,000	2.5										
2007	6,010	11,400	30,000	2.6	10,000 -									
2008	7,507	13,390	30,000	2.2										
2009	5,089	9,474	30,000	3.2	8,000 -								_	
2010	5,470	9,233	27,844	3.0		_					_	_		
2011	6,021	9,508	30,000	3.2	6,000 -									Ц
2012	6,978	11,086	33,000	3.0	,,,,,,			_						
2013	6,835	10,760	31,000	2.9	4.000									
2014	5,766	8,788	30,000	3.4	4,000									П
2015	6,895	7,671	29,000	3.8										
2016	7,948	8,243	29,000	3.5	2,000 -		Н		$\blacksquare$	▐	Н	-	Н	Н
2017	8,300	13,047	35,000	2.7										
2018	7,971	15,765	37,000	2.4				<b>.</b>						Ļ
2019	8,658	12,738	35,000	2.8		100 700 TO	90° 108°	100° 15	30,01	\$	3 %	50 50	\$ \$	2020
2020	11,671	17,758	41,000	2.3	1	, », »,	γ γ <sup>ν</sup>	かか	γ γ ·	マダ	, δ <sub>2</sub>	v v	γ,	V

RED=7 day season

POST-HUNT Deer Population	Trend	Stable
Vear Total		


Year	Total
2008	52,810
2009	58,926
2010	44,469
2011	46,503
2012	51,697
2013	55,713
2014	53,799
2015	47,438
2016	71,669
2017	74,053
2018	81,376
2019	60,756
2020	81,659
2021	95,277



#### DEER HEALTH: Fawn to Doe Ratio<sup>2</sup>

**Trend** Declining

Year	Total
2008	0.41
2009	0.43
2010	0.44
2011	0.40
2012	0.37
2013	0.38
2014	0.39
2015	0.40
2016	0.37
2017	0.39
2018	0.35
2019	0.33
2020	0.31




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

Regeneration Assessment Fa

Fair

Year % Adequate 2003-07 41% 2004-08 46% 2005-09 48% 2006-10 54% 2007-11 57% 2008-12 60% 2009-13 55% 2010-14 56% 2011-15 53% 2012-16 48% 2013-17 49% 2014-18 51% 2015-19 50%



Citizen Survey Results 2019 (2011)

**Too High** 24% (1

24% (11%) Just Right

47% (56%) **Too Low** 

23% (26%)

Antlerless Allocation Options					
Increase Stable Decr					
Firearm Season Option	Harvest	Harvest	Harvest		
7 day antlered & 7 day concurrent	43,000	38,000	33,000		
14 day concurrent	36,000	32,000	28,000		

<sup>&</sup>lt;sup>1</sup> - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

#### 2021-22 Pennsylvania Game Commission Antlerless Allocation Worksheet


			VVIV	io characteri	Stics	
WMU	2A	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		7%	61%	29%	3%	1.811

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	13,000
2005	8,510	19,649	55,000	2.7	12,000
2006	8,104	16,987	55,000	3.2	
2007	6,560	14,322	60,000	3.9	10,000
2008	6,714	15,255	55,000	3.5	
2009	6,829	13,920	55,000	4.0	8,000
2010	5,830	13,463	54,879	4.1	
2011	7,142	12,677	65,000	4.4	6,000
2012	6,683	12,694	59,000	4.5	
2013	6,836	13,241	49,000	3.7	
2014	5,131	9,580	46,000	4.8	4,000
2015	6,511	10,507	43,000	4.1	
2016	7,027	9,235	43,000	4.6	2,000
2017	6,134	10,866	50,000	4.6	
2018	6,036	10,950	49,000	4.2	_ <del>  1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,1,</del>
2019	6,929	9,918	46,000	4.4	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$
2020	8,128	11,835	46,000	3.9	

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	45,462
2009	50,336
2010	56,286
2011	49,033
2012	68,080
2013	53,996
2014	43,379
2015	30,033
2016	48,723
2017	57,963
2018	46,361
2019	44,587
2020	61,486
2021	72,156




#### DEER HEALTH: Fawn to Doe Ratio<sup>2</sup>

**Trend** 

Stable

Year	Total
2008	0.40
2009	0.38
2010	0.36
2011	0.39
2012	0.38
2013	0.39
2014	0.38
2015	0.31
2016	0.36
2017	0.39
2018	0.32
2019	0.32
2020	0.32




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

**Regeneration Assessment** Poor

% Adequate
46%
47%
46%
45%
44%
42%
43%
39%
43%
41%
41%
31%
28%



2019 (2011) **Citizen Survey Results** 

28% (25%) Just Right 50% (56%) **Too Low Too High** 19% (13%)

Antlerless Allocation Options					
Increase Stable Decr					
Firearm Season Option	Harvest	Harvest	Harvest		
7 day antlered & 7 day concurrent	53,000	45,000	38,000		
14 day concurrent	46,000	39,000	33,000		

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.

# 2021-22 Pennsylvania Game Commission Antlerless Allocation Worksheet

	WIVIU Characteristics				STICS	
WMU	2B	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		30%	44%	21%	0%	1,363

						Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12,000 —	
2005	5,182	14,459	68,000	4.4	12,000 —	
2006	5,759	16,505	68,000	3.9		
2007	4,372	15,332	68,000	3.9	10,000	
2008	3,964	15,251	68,000	4.1		
2009	4,297	19,866	68,000	3.3	8,000	
2010	3,976	13,008	68,000	4.8		
2011	4,472	16,550	71,000	3.6	6,000	
2012	4,837	15,955	67,000	3.8	1,000	
2013	5,610	14,389	62,000	4.3		-   -
2014	4,267	13,165	60,000	4.5	4,000	
2015	5,191	15,379	61,000	3.9		
2016	5,801	14,317	60,000	4.2	2,000 +	<del></del>
2017	4,458	13,930	60,000	3.9		
2018	5,036	12,318	58,000	3.8		<u>▊、▊、▊、▊、▊、▊、▊、▊、▋、▊、▊、▊、▊、▊、▊、</u>
2019	5,503	10,374	54,000	4.3	\\ \signiferance\s	z to
2020	6,201	14,746	49,000	3.3	1 20	\$ \$ \$ \$ \$ \$ \$ \$. \$ \$. \$ \$. \$ \$. \$ \$. \$

POST-HUNT Deer Population	Trend

Year Total


 $Harvest\ indices\ (i.e.,\ antlered\ harvest,\ antleress\ lic/deer),\ not\ PASAK\ model,\ used\ to\ monitor\ population\ trend$ 


Stable

#### **DEER HEALTH: Fawn to Doe Ratio<sup>2</sup>**

0.50					
0.45					
0.40 -		_			

**Trend** 





FOREST HEALTH			Regeneration Assessment
POREST HEALTH	Plot - Plot Regeneration	Plot - Plot Deer Impact	Mean Deer Impact

% Adequate Year

#### Forest data not considered in this developed WMU

Citizen Surve	ey Results	2019 (2011)		Too High	38%(32%)	Just Right	51% (52%)	Too Low	8% (9%)
Antlerless Allocation Options						-			
			Increses	Chabla	Dannaga	•			

Antieriess Allocation Options								
	Increase	Stable	Decrease					
Firearm Season Option	Harvest	Harvest	Harvest					
14 day concurrent	53,000	49,000	42,000					

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

#### 2021-22 Pennsylvania Game Commission Antlerless Allocation Worksheet


			VVIV	WIVIO Characteristics					
WMU	2C	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)			
		6%	68%	24%	10%	2.934			

									Antle	red I	larve	est						
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12,000													
2005	7,413	13,683	53,000	3.8	12,000 -													
2006	9,049	12,094	49,000	4.0														
2007	8,441	11,619	49,000	4.1	10,000 -													_
2008	7,476	12,752	49,000	3.8													ı	
2009	6,508	10,870	49,000	4.5	8,000 -		-				<del></del>			Н			-	-
2010	8,528	9,579	44,107	4.6													ı	ı
2011	8,249	12,793	58,000	4.5	6,000 -	H					Ш	┸		Ш			4	
2012	7,600	10,822	50,000	4.6	,,,,,,												ı	ı
2013	7,219	10,957	43,000	3.9													ı	ı
2014	7,016	8,985	38,000	4.5	4,000 -	П											ı	1
2015	9,134	7,269	31,000	4.3													ı	
2016	8,300	6,869	31,000	4.6	2,000 -			-			H	╂	$\blacksquare$	Н			╅	-
2017	9,792	7,724	31,000	4.0													ı	
2018	9,572	11,134	44,000	4.0			_		┍┻┦		Ļ∎	<b>,</b> ■				_		_
2019	9,426	12,743	52,000	4.1		ip <sup>J</sup> ap	2001	200° 1	00° 00	\$ \$	3	\$	2 <sup>th</sup> 203	;	3	2029	1019 Y	B
2020	8,441	15,744	58,000	3.7	1 1	, \$	2	20° 1	> 5.	\$	\$ .	b y	5 V	20.	20.	₽.	₽`^	ν.

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
Teal	TOLAI
2008	87,046
2009	72,402
2010	62,340
2011	66,729
2012	64,888
2013	61,386
2014	68,683
2015	66,027
2016	83,350
2017	69,034
2018	113,659
2019	86,087
2020	97,246
2021	76,365




#### DEER HEALTH: Fawn to Doe Ratio<sup>2</sup>

Approximately 50% of WMU is a CWD DMA in 2019

**Trend** 

Stable

Year	Total
2008	0.43
2009	0.43
2010	0.40
2011	0.40
2012	0.38
2013	0.44
2014	0.42
2015	0.37
2016	0.35
2017	0.37
2018	0.34
2019	0.35
2020	0.33




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change Mean Deer Impact

Fair

2 1

Year	% Adequate
2003-07	54%
2004-08	59%
2005-09	58%
2006-10	58%
2007-11	59%
2008-12	56%
2009-13	57%
2010-14	58%
2011-15	62%
2012-16	63%
2013-17	60%
2014-18	58%
2015-19	57%



Citizen Survey Results 2019 (2011)

Too High

19%(13%) Just Right

52%(50%) **Too Low** 

23%(26%)

Antlerless Allocation Options										
	Increase	Stable	Decrease							
Firearm Season Option	Harvest	Harvest	Harvest							
14 day concurrent	67,000	51,000	40,000							

<sup>&</sup>lt;sup>1</sup> - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

#### 2021-22 Pennsylvania Game Commission Antlerless Allocation Worksheet


			VVIV	WIVIO Characteristics					
WMU	2D	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)			
		5%	60%	31%	2%	2.486			

Deer Harvest						
					Antlered Harvest	
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	14.000	
2005	9,975	22,054	56,000	2.5	14,000	
2006	10,896	20,437	56,000	2.7		
2007	9,118	18,099	56,000	3.1	12,000	
2008	9,508	15,591	56,000	3.5		
2009	9,977	15,962	56,000	3.5	10,000	
2010	11,540	18,046	50,123	2.8		
2011	11,130	19,257	60,000	3.1	8,000	
2012	13,660	20,839	62,000	3.0		
2013	13,704	21,614	61,000	2.8		
2014	11,417	16,441	61,000	3.7	6,000	
2015	12,292	15,728	55,000	3.5		
2016	12,843	16,447	55,000	3.3	4,000	
2017	14,716	17,033	55,000	3.2		
2018	11,847	20,345	63,000	3.1	2,000	
2019	12,971	17,472	66,000	3.8	\$	
2020	12,121	18,726	60,000	3.2		

RED=7-day season

	POST-HUNT Deer Population	Trend	Stable
--	---------------------------	-------	--------

Year	Total
2008	69,732
2009	88,666
2010	86,493
2011	101,182
2012	102,440
2013	113,774
2014	144,084
2015	110,214
2016	117,823
2017	112,499
2018	140,281
2019	105,280
2020	114,679
2021	93,498




#### DEER HEALTH: Fawn to Doe Ratio<sup>2</sup>

Approximately 13% of WMU is a CWD DMA in 2019

**Trend** 

Declining

Year	Total
2008	0.44
2009	0.39
2010	0.42
2011	0.44
2012	0.42
2013	0.42
2014	0.40
2015	0.37
2016	0.36
2017	0.37
2018	0.34
2019	0.33
2020	0.32



**FOREST HEALTH** 


Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

Fair

**Regeneration Assessment** 

Mean Deer Impact 3 or less

Year	% Adequate
2003-07	52%
2004-08	54%
2005-09	51%
2006-10	52%
2007-11	49%
2008-12	46%
2009-13	50%
2010-14	45%
2011-15	44%
2012-16	50%
2013-17	48%
2014-18	41%
2015-19	45%



**Citizen Survey Results** 2019 (2011)

26%(23%) Just Right

57%(52%) Too Low

13%(19%)

Antlerless Allocation Options						
	Increase	Stable	Decrease			
Firearm Season Option	Harvest	Harvest	Harvest			
14 day concurrent	74,000	62,000	55,000			

<sup>&</sup>lt;sup>1</sup> - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

#### 2021-22 Pennsylvania Game Commission Antlerless Allocation Worksheet


			VVIV	io characteri	Stics	
WMU	2E	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		5%	65%	26%	6%	1.427

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12,000
2005	4,093	7,471	21,000	2.8	12,000
2006	5,358	7,360	21,000	2.8	
2007	3,642	6,398	21,000	3.2	10,000
2008	4,984	6,179	21,000	3.3	
2009	3,673	5,298	21,000	4.0	8,000
2010	4,178	5,952	20,407	3.5	_
2011	4,116	7,073	25,000	3.5	6,000
2012	4,785	5,561	21,000	3.8	
2013	4,883	7,973	22,000	2.8	
2014	4,440	5,593	21,000	3.8	4,000
2015	4,742	5,263	21,000	4.0	
2016	5,221	5,215	21,000	4.1	2,000
2017	6,929	6,214	22,000	3.5	
2018	6,274	8,693	27,000	3.1	_ <u> </u>
2019	6,370	7,641	32,000	4.2	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$
2020	6,515	11,348	39,000	3.4	\(\rightarrow\rightarr

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	32,623
2009	42,709
2010	38,317
2011	38,134
2012	30,384
2013	44,546
2014	45,529
2015	50,549
2016	43,081
2017	43,144
2018	56,635
2019	47,171
2020	62,753
2021	52,578




Approximately 51% of WMU is a CWD DMA in 2019

**Trend** 

Stable

Year	Total
2008	0.40
2009	0.37
2010	0.46
2011	0.41
2012	0.43
2013	0.40
2014	0.36
2015	0.36
2016	0.36
2017	0.33
2018	0.31
2019	0.33
2020	0.32



**FOREST HEALTH** 


Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

Fair

**Regeneration Assessment** 

Mean Deer Impact 3 or less

Year	% Adequate
2003-07	53%
2004-08	50%
2005-09	47%
2006-10	50%
2007-11	52%
2008-12	52%
2009-13	56%
2010-14	61%
2011-15	63%
2012-16	56%
2013-17	60%
2014-18	56%
2015-19	54%



**Citizen Survey Results** 2019 (2011)

20%(13%) Just Right

56%(48%) **Too Low** 

22%(31%)

Antlerless Allocation Options						
Increase Stable Decrea						
Firearm Season Option	Harvest	Harvest	Harvest			
14 day concurrent	42,000	35,000	30,000			

<sup>1 -</sup> The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.


			WIVIO Characteristics				
WMU	2F	% Develope	d % Forest	%Ag/Field	%Public	Area (sq mi)	
		2%	88%	7%	56%	2.409	

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	13,000
2005	6,013	8,322	30,000	3.5	12,000
2006	7,153	8,030	28,000	3.5	, <u>, , , , , , , , , , , , , , , , , , </u>
2007	4,795	7,132	28,000	3.9	10,000
2008	6,990	9,117	28,000	3.0	
2009	5,167	6,648	28,000	4.3	8,000
2010	6,403	5,657	22,148	4.0	
2011	5,393	6,737	34,000	5.0	6,000
2012	7,139	6,067	27,000	4.5	
2013	6,607	8,008	29,000	3.6	
2014	5,979	5,915	27,000	4.6	4,000
2015	6,989	5,434	22,000	4.1	
2016	7,678	6,718	22,000	3.3	2,000
2017	9,489	7,200	24,000	3.3	
2018	7,665	7,533	23,000	3.1	_
2019	9,014	8,816	31,000	3.5	\$
2020	10,686	9,953	36,000	3.6	

RED=7-day season

POST-HUNT Deer Population	Trend	Increasing

Year	Total
2008	47,288
2009	67,724
2010	46,887
2011	70,765
2012	53,210
2013	83,063
2014	65,614
2015	61,020
2016	67,152
2017	74,387
2018	108,575
2019	87,309
2020	98,104
2021	112,840




Approximately 8% of WMU is a CWD DMA in 2019

**Trend** 

Stable

Year	Total
2008	0.35
2009	0.34
2010	0.39
2011	0.40
2012	0.39
2013	0.38
2014	0.37
2015	0.34
2016	0.33
2017	0.36
2018	0.37
2019	0.28
2020	0.27




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

Regeneration Assessment Good

Good

Year % Adequate 2003-07 47% 2004-08 50% 2005-09 50% 2006-10 54% 2007-11 54% 2008-12 54% 2009-13 54% 2010-14 58% 2011-15 61% 2012-16 65% 2013-17 69% 2014-18 71% 2015-19 69%



Citizen Survey Results 2019 (2011)

Too High

19%(10%) Just Right

48%(39%) **Too Low** 

26%(42%)

Antlerless Allocation Options							
Increase Stable Decreas							
Firearm Season Option	Harvest	Harvest	Harvest				
7 day antlered & 7 day concurrent	40,000	32,000	23,000				
14 day concurrent	32,000	25,000	18,000				

<sup>&</sup>lt;sup>1</sup> - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{2}</sup>$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			WIVIO Characteristics				
WMU	2G	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)	
		4%	82%	7%	57%	3.117	

					Antlered Harvest	
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	13.000	_
2005					12,000	
2006						
2007					10,000	_
2008						
2009	3,802	1,046			8,000	_
2010	5,088	2,627				
2011	4,957	4,117			6,000	_
2012	4,976	4,915				
2013	5,018	6,881	28,000	4.1		
2014	4,839	4,671	22,000	4.7	4,000	
2015	6,073	4,143	22,000	5.4		
2016	6,201	3,996	21,000	5.3	2,000	1
2017	8,193	5,516	25,500	4.6		
2018	6,296	7,372	30,000	4.1		_
2019	8,062	6,123	26,000	4.3	50 50 50 50 50 50 50, 50, 50, 50, 50, 50	SP
2020	7,505	6,806	27,000	4.0		>

RED=7-day season


Deer Population	Trend	Stable

Year	Total
2008	
2009	55,234
2010	41,008
2011	45,743
2012	55,997
2013	57,014
2014	49,313
2015	40,343
2016	65,521
2017	67,942
2018	81,757
2019	55,221
2020	70,946
2021	85,558



Trend Stable

Year	Total
2008	0.34
2009	0.35
2010	0.39
2011	0.35
2012	0.25
2013	0.35
2014	0.30
2015	0.34
2016	0.31
2017	0.29
2018	0.35
2019	0.26
2020	0.24




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

Regeneration Assessment Fa

Fair

Year	% Adequate
2003-07	53%
2004-08	53%
2005-09	54%
2006-10	54%
2007-11	55%
2008-12	56%
2009-13	55%
2010-14	54%
2011-15	56%
2012-16	52%
2013-17	55%
2014-18	52%
2015-19	50%



Citizen Survey Resul	ts 2019	(2011)
----------------------	---------	--------

**Too High** 13%(3%) **Just Right** 49%(39%) **Too Low** 35%(55%)

Antlerless Allocation Options				
Increase Stable Decrease				
Firearm Season Option	Harvest	Harvest	Harvest	
7 day antlered & 7 day concurrent	41,000	28,000	15,000	
14 day concurrent	34,000	23,000	12,000	

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.

			WMU Characteristics			
WMU	2H	% Deve	loped % Forest	%Ag/Field	%Public	Area (sq mi)
		49	6 86%	6%	27%	1.001

er Harve	est					
					Antlered Harve	est
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12.000	
2005					12,000	
2006						
2007					10,000	
2008						
2009	1,471	1,046			8,000	
2010	1,670	990				
2011	1,323	1,321			6,000	
2012	1,565	1,459			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
2013	1,475	1,657	6,000	3.7		
2014	1,670	1,064	5,500	5.2	4,000	
2015	1,426	1,419	6,500	4.6		1
2016	1,867	1,861	6,000	3.2	2,000	
2017	1,726	1,889	7,000	3.7		
2018	2,478	1,812	6,000	3.3	- <del>                                    </del>	<del>, , , , , , , , , , , , , , , , , , , </del>
2019	2,404	1,086	6,000	5.6	Lag Lag Lag Lag Lag Lag Lag Lag.	344444
2020	2,855	1,563	7,000	4.5	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	× × × × × × × × × × × × × × × × × × ×


RED=7-day season

eer Popul	ation		Trend Increasing
Year	Total		
2008		160,000	
2009	19,730	140,000	
2010	11,565	-13,113	
2011	18,952	120,000	
2012	13,917		
2013	16,895	100,000	
2014	16,537	80,000	
2015	16,872	50,000	
2016	15,430	60,000	
2017	15,704		
2018	38,649	40,000	
2019	18,919	20,000	
2020	25,314	· ·	
2021	42,858	0	<u>┛╷┛╷┛╷┛╷┛╷┚╷┚╷┚</u> ╷
		Les Les Les Les Les	

**Trend** 

Stable

Year	Total
2008	0.43
2009	0.40
2010	0.35
2011	0.33
2012	0.39
2013	0.23
2014	0.33
2015	0.33
2016	0.24
2017	0.40
2018	0.35
2019	0.32
2020	0.22




**FOREST HEALTH** 

**Regeneration Assessment** Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

Fair

Year	% Adequate
2003-07	44%
2004-08	43%
2005-09	47%
2006-10	52%
2007-11	51%
2008-12	51%
2009-13	57%
2010-14	56%
2011-15	57%
2012-16	55%
2013-17	62%
2014-18	54%
2015-19	50%



Citizen	Survey	Results	2019	(2011)	i

**Too High** 

13%(3%) Just Right

49%(39%) Too Low

35%(55%)

Antlerless Allocation Options				
Firearm Season Option	Increase Harvest	Stable Harvest	Decrease Harvest	
7 day antlered & 7 day concurrent	11,000	7,000	2,000	
14 day concurrent	9,000	6,000	2,000	

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.

			VVIV	WIVIO Characteristics		
WMU	3A	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		2%	78%	17%	10%	1.506

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12.000
2005	3,981	8,657	27,000	3.1	12,000
2006	4,527	8,818	29,000	3.2	
2007	3,359	7,803	29,000	3.6	10,000
2008	4,132	7,478	26,000	3.4	
2009	3,310	5,998	26,000	4.4	8,000
2010	3,751	6,469	25,247	3.9	
2011	3,345	6,672	26,000	3.9	6,000
2012	4,278	6,673	26,000	3.9	
2013	4,177	5,430	23,000	4.2	
2014	3,308	4,253	18,000	4.2	4,000
2015	4,314	4,005	19,000	4.8	
2016	5,432	3,776	15,000	4.0	2,000
2017	5,419	5,014	20,000	4.0	
2018	4,825	7,430	22,000	3.0	_ <del>  _  <b>                               </b></del>
2019	5,704	5,663	20,000	3.5	\$
2020	6,968	6,694	21,000	3.1	


RED=7-day season

ST-HUNT	Deer Population		Trend	Stable
Year	Total	160,000		
2008	32,425	100,000		
2009	32,513	140,000		
2010	31,412			
2011	39,532	120,000		
2012	31,224	100,000		
2013	41,358	100,000		
2014	45,317	80,000		
2015	36,181			
2016	49,307	60,000		
2017	49,426	40.000		
2018	55,441	40,000		
2019	39,832	20,000		
2020	54,040			
2021	71,376	0 +		

Trend

Stable

Year	Total
2008	0.33
2009	0.35
2010	0.42
2011	0.36
2012	0.34
2013	0.36
2014	0.36
2015	0.36
2016	0.27
2017	0.32
2018	0.33
2019	0.32
2020	0.26




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

Regeneration Assessment Fair

Fair

% Adequate
65%
63%
62%
61%
63%
60%
66%
66%
67%
65%
69%
64%
61%



Citizen Survey Resul	ts 2019	(2011)
----------------------	---------	--------

**Too High** 18(3%) **Just Right** 57%(32%) **Too Low** 21%(59%)

Antlerless Allocation Options								
Firearm Season Option	Increase Harvest	Stable Harvest	Decrease Harvest					
7 day antlered & 7 day concurrent	26,000	21,000	16,000					
14 day concurrent	24,000	19,000	15,000					

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.


			io characteri	Stics		
WMU	3B	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		6%	79%	11%	21%	2.218

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12.000
2005	5,980	10,871	41,000	3.7	12,000
2006	6,530	10,563	43,000	4.0	
2007	5,933	10,177	43,000	4.2	10,000
2008	5,469	9,857	43,000	4.3	
2009	4,865	9,112	43,000	4.7	8,000
2010	5,369	7,585	33,761	4.5	_ [ ] _ [
2011	5,935	7,707	40,000	5.2	6,000
2012	5,752	8,701	40,000	4.6	
2013	6,153	8,718	39,000	4.5	
2014	6,039	8,055	33,000	4.1	4,000
2015	6,840	7,359	28,000	3.8	
2016	7,481	7,290	28,000	3.8	2,000
2017	8,945	6,970	30,000	4.3	
2018	6,977	8,354	29,000	3.5	<u> </u>
2019	7,558	10,264	38,000	3.7	6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6
2020	9,090	8,507	33,000	3.9	

RED=7-day season

Deer Population	Trend	Stable
-----------------	-------	--------


Year	Total
2008	56,162
2009	46,869
2010	48,895
2011	49,768
2012	58,481
2013	53,709
2014	63,803
2015	55,249
2016	76,808
2017	80,598
2018	76,249
2019	51,976
2020	62,489
2021	90,795



T	re	nc	

Stable

Year	Total
2008	0.37
2009	0.40
2010	0.40
2011	0.38
2012	0.39
2013	0.39
2014	0.38
2015	0.38
2016	0.37
2017	0.33
2018	0.31
2019	0.30
2020	0.35




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

**Regeneration Assessment** Good

17%(24%)

Year	% Adequate
2003-07	58%
2004-08	59%
2005-09	62%
2006-10	62%
2007-11	60%
2008-12	65%
2009-13	67%
2010-14	65%
2011-15	61%
2012-16	64%
2013-17	57%
2014-18	63%
2015-19	66%



Citizen Survey Resul	ts 2019	(2011)
----------------------	---------	--------

Too High 20%(7%) Just Right 55%(59%) **Too Low** 

Antlerless Allocation Options						
Increase Stable Decrease						
Firearm Season Option	Harvest	Harvest	Harvest			
7 day antlered & 7 day concurrent	42,000	33,000	25,000			
14 day concurrent	39,000	30,000	23,000			

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.


			VVIV	io characteri	Stics	
WMU	3C	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		4%	75%	16%	3%	2.187

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12,000
2005	5,821	11,198	32,000	2.8	12,000
2006	6,673	9,248	27,000	2.9	
2007	5,278	9,586	27,000	2.8	10,000
2008	6,288	7,258	27,000	3.7	
2009	6,196	7,084	27,000	3.9	8,000
2010	6,211	8,309	26,358	3.2	
2011	7,103	9,943	29,000	2.9	6,000
2012	7,854	10,508	35,000	3.3	
2013	7,004	12,683	35,000	2.8	
2014	6,526	10,302	32,000	3.1	4,000
2015	7,614	10,460	36,000	3.4	
2016	8,629	10,968	36,000	3.3	2,000
2017	8,703	11,860	42,000	3.5	
2018	7,739	12,172	38,000	3.1	<u> </u>
2019	9,382	12,808	46,000	3.6	\$ \$
2020	10,843	14,538	49,000	3.4	

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

45,511 54,141 65,624
65,624
,
EO 34E
59,245
64,359
67,720
58,925
67,997
83,206
85,083
79,925
57,169
75,360
94,807



Declining

#### DEER HEALTH: Fawn to Doe Ratio<sup>2</sup>

**Total** 

0.31

0.35

0.32

0.34

0.32

0.35

0.36

0.34

0.38

0.33

0.32

0.27

0.29

Year

2008

2009

2010

2011

2012

2013

2014

2015

2016

2017

2018

2019

2020

0.50	
0.45	
0.40	
0.35	
0.30	<del>                                      </del>
0.25	<del></del>
0.20	<del> </del>
0.15	
0.10	<del></del>
0.05	. <del></del>
0.00	<b></b>
708	* 68 60 62 62 62 62 62 62 62 62 62 62

**Trend** 


**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

**Regeneration Assessment** Mean Deer Impact 3 or less

Fair

Year	% Adequate
2003-07	49%
2004-08	53%
2005-09	53%
2006-10	51%
2007-11	51%
2008-12	54%
2009-13	56%
2010-14	55%
2011-15	53%
2012-16	53%
2013-17	50%
2014-18	44%
2015-19	46%



**Citizen Survey Results** 2019 (2011) Too High

30%(10%) Just Right

55%(61%) **Too Low** 

11%(20%)

Antlerless Allocation Options						
Increase Stable Decre Firearm Season Option Harvest Harvest Harv						
7 day antlered & 7 day concurrent	52,000	44,000	37,000			
14 day concurrent	39,000	33,000	28,000			

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			WIV	/IU Characteri	stics	
WMU	3D	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		11%	7/1%	6%	16%	2 101

er Harve						Antlered Harvest	_
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12.000		_
2005	3,865	7,254	38,000	5.1	12,000 —		
2006	4,969	7,445	38,000	5.0			
2007	3,647	7,017	38,000	5.3	10,000 +		_
2008	3,899	6,925	37,000	5.3			
2009	3,096	6,265	37,000	5.9	8,000		_
2010	3,884	5,509	31,622	5.8			
2011	4,509	7,163	39,000	5.4	6,000		_
2012	4,039	6,010	39,000	6.5	0,000	_	ı
2013	3,446	4,986	32,000	6.4		I	ı
2014	4,155	5,203	25,000	4.8	4,000		
2015	3,500	3,655	25,000	6.9			ı
2016	4,272	4,235	25,000	5.9	2,000 +	<del> </del>	┢
2017	4,656	4,187	25,000	5.9			
2018	5,189	5,690	25,000	4.4	- +	<del>┖╷▊╷▊╷▊╷▊╷▊╷▊╷▊╷▊╷▊</del> ╷▊ <del>╷</del> ▊	Ļ
2019	6,016	4,932	25,000	5.1	\\ \delta^6	be to	, S
2020	6,180	6,366	36,000	5.7	1 %	****	∾

RED=7-day season

POST-HUNT Deer Population	Trend	Increasing


Year	Total
2008	31,623
2009	37,563
2010	25,378
2011	30,250
2012	31,299
2013	29,225
2014	25,127
2015	33,778
2016	28,957
2017	33,302
2018	30,727
2019	33,798
2020	48,663
2021	45,355



**Trend** 

Stable

Year	Total
2008	0.32
2009	0.35
2010	0.31
2011	0.34
2012	0.34
2013	0.30
2014	0.29
2015	0.27
2016	0.31
2017	0.24
2018	0.30
2019	0.27
2020	0.29




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

**Regeneration Assessment** Fair

>3

Year	% Adequate
2003-07	56%
2004-08	54%
2005-09	55%
2006-10	58%
2007-11	57%
2008-12	59%
2009-13	61%
2010-14	61%
2011-15	57%
2012-16	63%
2013-17	57%
2014-18	59%
2015-19	58%



2019 (2011) **Citizen Survey Results** 

30%(13%) Just Right **Too High** 52%(57%) **Too Low** 13%(24%)

Antlerless Allocation Options				
Firearm Season Option	Increase Harvest	Stable Harvest	Decrease Harvest	
7 day antlered & 7 day concurrent	39,000	29,000	18,000	
14 day concurrent	36,000	27,000	17,000	

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.


			VVIV	WIVIO Characteristics		
WMU	4A	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		4%	70%	24%	15%	1.736

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	42.000
2005	3,714	7,578	35,000	4.5	12,000
2006	5,871	7,827	29,000	3.6	
2007	4,477	6,735	29,000	4.2	10,000
2008	4,187	6,874	29,000	4.2	
2009	3,733	7,414	29,000	3.9	8,000
2010	3,761	6,401	27,521	4.3	
2011	4,849	6,527	28,000	4.3	6,000
2012	4,245	6,463	29,000	4.5	
2013	4,961	5,981	28,000	4.7	
2014	3,317	6,802	28,000	5.6	4,000
2015	5,095	6,360	30,000	4.7	
2016	4,423	5,726	30,000	5.2	2,000
2017	4,810	6,475	30,000	4.6	
2018	5,142	6,395	38,000	5.5	<u> </u>
2019	5,981	5,250	41,000	5.8	\$ \$
2020	5,183	10,849	49,000	4.0	

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	47,414
2009	34,628
2010	30,789
2011	38,125
2012	49,191
2013	36,579
2014	42,196
2015	23,772
2016	48,538
2017	29,746
2018	39,238
2019	42,174
2020	47,047
2021	39,911




100% of WMU is a CWD DMA in 2019

**Trend** 

Stable

Year	Total
2008	0.34
2009	0.34
2010	0.32
2011	0.38
2012	0.32
2013	0.30
2014	0.32
2015	0.37
2016	0.29
2017	0.30
2018	0.30
2019	0.30
2020	0.29




**FOREST HEALTH** 

**Regeneration Assessment** Good

3 or less

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change Mean Deer Impact

Year % Adequate 2003-07 60% 2004-08 64% 2005-09 64% 2006-10 61% 2007-11 63% 2008-12 60% 2009-13 59% 2010-14 61% 2011-15 63% 2012-16 67% 2013-17 68% 2014-18 75% 2015-19 67%



Citizen Survey Results 2019 (2011)

**Too High** 

14%(4%) Just Right

45%(45%) Too Low

37%(42%)

Antlerless Allocation Options	Antlerless	Allocation	<b>Options</b>
-------------------------------	------------	------------	----------------

	Increase	Stable	Decrease
Firearm Season Option	Harvest	Harvest	Harvest
14 day concurrent	50.000	36.000	29.000

<sup>&</sup>lt;sup>1</sup> - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.


			WIVIO Characteristics				
WMU	4B	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)	
		6%	65%	27%	15%	1.591	

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12.000
2005	3,571	6,644	35,000	5.2	12,000
2006	5,026	6,626	31,000	4.6	
2007	3,472	4,509	23,000	5.0	10,000
2008	3,917	3,846	23,000	5.9	
2009	4,011	4,061	23,000	5.7	8,000
2010	4,458	5,113	22,148	4.4	
2011	5,341	5,498	23,000	4.2	6,000
2012	5,622	5,636	26,000	4.6	
2013	5,312	5,769	24,000	4.2	
2014	4,611	5,630	26,000	4.6	4,000
2015	5,701	6,961	26,000	3.8	
2016	5,164	6,151	26,000	4.2	2,000
2017	5,602	7,061	26,000	3.7	
2018	5,273	6,757	26,000	3.9	<u> </u>
2019	5,722	7,305	32,000	4.4	\$ \$
2020	5,034	10,770	33,000	3.1	

RED=7-day season

POST-HUNT Deer Population	Trend	Decreasing

Year	Total
2008	30,479
2009	39,044
2010	43,550
2011	37,273
2012	60,340
2013	52,903
2014	50,517
2015	45,362
2016	57,846
2017	55,941
2018	52,407
2019	50,252
2020	54,044
2021	44,691




DEER HEALTH: Fawn to Doe Ratio<sup>2</sup> 100% of WMU is a CWD DMA in 2019

Trend

Declining

Year	Total
2008	0.38
2009	0.44
2010	0.38
2011	0.41
2012	0.33
2013	0.31
2014	0.40
2015	0.36
2016	0.35
2017	0.34
2018	0.30
2019	0.29
2020	0.32




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change Mean Deer Impact

Fair

I all

Year	% Adequate
2003-07	61%
2004-08	60%
2005-09	58%
2006-10	60%
2007-11	64%
2008-12	61%
2009-13	59%
2010-14	60%
2011-15	63%
2012-16	68%
2013-17	59%
2014-18	57%
2015-19	58%



Citizen Survey Results 2019 (2011)
------------------------------------

**Too High** 16%(6%) **Just Right** 53%(53%) **Too Low** 21%(33%)

Antlerless	Allocation Opti	ons	
	Increase	Stable	Decrease
Firearm Season Option	Harvest	Harvest	Harvest
14 day concurrent	34,000	27,000	22,000

<sup>&</sup>lt;sup>1</sup> - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	io Characteri	Stics	
WMU	4C	% Developed	d % Forest	%Ag/Field	%Public	Area (sq mi)
		8%	71%	17%	15%	1.717

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12.000
2005	5,891	9,805	39,000	3.9	12,000
2006	6,115	8,883	39,000	4.2	
2007	4,828	9,375	39,000	4.1	10,000
2008	5,015	8,027	35,000	4.3	
2009	4,745	7,163	35,000	4.9	8,000
2010	5,724	8,357	34,351	4.2	
2011	5,525	7,392	35,000	4.7	6,000
2012	5,335	7,823	35,000	4.5	
2013	5,180	6,922	27,000	3.9	
2014	4,830	4,996	25,000	5.1	4,000
2015	5,381	4,976	25,000	5.1	
2016	6,381	5,273	25,000	4.8	2,000
2017	6,799	6,464	29,000	4.5	
2018	5,781	7,155	30,000	4.2	_ <del>  <b>                                  </b></del>
2019	6,975	8,328	36,000	4.3	\$ \$
2020	6,998	8,055	32,000	4.0	

RED=7-day season

POST-HUNT Deer Population	Trend	Stable


Year	Total
2008	44,569
2009	45,224
2010	44,256
2011	58,091
2012	45,093
2013	45,586
2014	49,072
2015	50,265
2016	55,068
2017	55,311
2018	61,317
2019	55,122
2020	55,238
2021	77,639



**Trend** 

Stable

Year	Total
2008	0.42
2009	0.41
2010	0.42
2011	0.38
2012	0.39
2013	0.40
2014	0.40
2015	0.37
2016	0.36
2017	0.37
2018	0.37
2019	0.33
2020	0.34




**FOREST HEALTH** 

**Regeneration Assessment** Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

Fair

Year	% Adequate
2003-07	66%
2004-08	63%
2005-09	63%
2006-10	63%
2007-11	60%
2008-12	61%
2009-13	62%
2010-14	58%
2011-15	60%
2012-16	59%
2013-17	60%
2014-18	61%
2015-19	59%



Citizen Survey F	Results 2019	(2011)	
------------------	--------------	--------	--

Too High 23%(7%)

**Just Right** 

52%(56%) **Too Low** 

21%(26%)

Antlerless Allocation Options				
	Increase	Stable	Decrease	
Firearm Season Option	Harvest	Harvest	Harvest	
7 day antlered & 7 day concurrent	40,000	33,000	26,000	
14 day concurrent	35,000	29,000	23,000	

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.


			VVIV	io characteri	STICS	
WMU	4D	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		6%	70%	22%	28%	2,743

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	12.000
2005	5,591	8,354	40,000	4.7	12,000
2006	6,776	9,878	40,000	4.0	_
2007	5,765	8,073	40,000	4.9	10,000
2008	6,593	9,310	40,000	4.2	_ [
2009	4,971	7,192	40,000	5.6	8,000
2010	6,321	5,472	30,052	5.6	
2011	7,144	6,561	37,000	5.7	6,000
2012	6,922	6,325	36,000	5.7	
2013	7,165	8,225	35,000	4.3	
2014	6,461	6,832	33,000	5.0	4,000
2015	7,240	7,197	33,000	4.6	
2016	7,921	7,234	34,000	4.7	2,000
2017	10,594	8,381	34,000	4.0	
2018	8,299	8,703	34,000	3.9	_ <del>                                     </del>
2019	8,740	10,266	46,000	4.5	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$
2020	9,141	12,256	45,000	3.7	

RED=7-day season


POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	43,299
2009	62,529
2010	46,284
2011	73,017
2012	70,495
2013	67,011
2014	61,428
2015	56,905
2016	60,398
2017	63,984
2018	99,997
2019	61,822
2020	71,983
2021	89,963




#### DEER HEALTH: Fawn to Doe Ratio<sup>2</sup> Approximately 30% of WMU is a CWD DMA in 2019 Trend Declining

Total
0.34
0.31
0.39
0.36
0.37
0.36
0.31
0.32
0.31
0.31
0.29
0.30
0.29



FOREST HEALTH Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change Mean Deer Impact 3 or less

% Adequate
43%
43%
43%
44%
43%
48%
49%
48%
52%
53%
48%
50%
52%



Citizen Survey Results 2019 (2011) Too High 20% (8%) Just I	Right 48%(46%) Too Low	26%(38%)
---	------------------------	----------

Antlerless	Allocation Opti	ons	
	Increase	Stable	Decrease
Firearm Season Option	Harvest	Harvest	Harvest
14 day concurrent	55,000	40,000	29,000

 $<sup>^{1}</sup>$  - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	io Characteri	Stics	
WMU	4E	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		8%	54%	34%	4%	1.736

						Α	ntlered	d Harve	est				
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	13.000								
2005	4,544	9,130	38,000	4.1	12,000								
2006	4,134	8,975	38,000	4.2									
2007	3,314	8,119	38,000	4.6	10,000								
2008	4,270	7,193	30,000	4.1									
2009	4,064	6,287	30,000	4.8	8,000							_	
2010	4,768	5,923	26,899	4.6								l .	
2011	5,076	6,054	29,000	4.8	6,000								
2012	4,960	6,079	28,000	4.6	5,555								
2013	6,287	7,707	26,000	3.4	4.000	_							
2014	5,847	5,919	21,000	3.6	4,000	_							
2015	6,202	6,914	25,000	3.6									
2016	7,294	7,474	25,000	3.4	2,000				$\blacksquare$			Н	-
2017	8,241	8,735	27,500	3.1									
2018	6,980	9,345	32,000	3.4	- <b>  II , II</b> ,								
2019	7,314	9,513	34,000	3.6	Log Logo L	io par for	2 20 C	5,00	03,04	, 50 50 C	\$ \$	103°	00° 60
2020	8,625	11,209	37,000	3.3	~ ~ ~	, 20 V	, 25 X	, 20 (	\$ \$	2 2	, <sub>2</sub>	γ ·	7 8

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	35,121
2009	37,339
2010	36,311
2011	51,706
2012	44,225
2013	48,318
2014	50,707
2015	59,206
2016	64,923
2017	62,285
2018	70,064
2019	60,055
2020	59,120
2021	77,399




Approximately 4% of WMU is a CWD DMA in 2019

**Trend** 

Stable

Vanu	Total
Year	Total
2008	0.44
2009	0.44
2010	0.44
2011	0.42
2012	0.45
2013	0.32
2014	0.38
2015	0.40
2016	0.35
2017	0.34
2018	0.33
2019	0.29
2020	0.34




**FOREST HEALTH** 

Plot - Plot Regeneration No Change Plot - Plot Deer Impact No Change

**Regeneration Assessment** Fair **Mean Deer Impact** 

>3

Year	% Adequate
2003-07	68%
2004-08	68%
2005-09	65%
2006-10	66%
2007-11	65%
2008-12	60%
2009-13	64%
2010-14	56%
2011-15	56%
2012-16	67%
2013-17	69%
2014-18	65%
2015-19	64%



**Citizen Survey Results** 2019 (2011) **Too High** 

30%(8%)

**Just Right** 

50%(58%) **Too Low** 

16%(28%)

Antlerless Allocation Options						
<b>.</b>	Increase	Stable	Decrease			
Firearm Season Option	Harvest	Harvest	Harvest			
7 day antlered & 7 day concurrent	43,000	35,000	29,000			
14 day concurrent	42,000	34,000	28,000			

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.


			VVIV	io Characteri	Stics	
WMU	5A	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		14%	35%	49%	11%	1.301

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	13,000
2005	2,396	4,690	28,000	5.8	12,000
2006	2,155	5,207	25,000	4.7	
2007	2,433	3,881	22,000	5.5	10,000
2008	2,057	3,778	19,000	4.9	
2009	2,237	4,194	19,000	4.6	8,000
2010	2,442	3,398	18,269	5.4	
2011	3,575	3,573	19,000	5.3	6,000
2012	2,795	3,596	19,000	5.3	3,555
2013	2,825	4,098	19,000	4.6	
2014	2,377	3,282	19,000	5.8	4,000
2015	2,862	4,631	19,000	4.1	
2016	3,017	4,047	19,000	4.7	2,000
2017	2,925	3,811	22,000	5.7	
2018	3,091	4,649	23,000	4.9	<u> </u>
2019	3,406	4,951	22,000	4.4	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$
2020	3,522	6,087	26,000	4.3	

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	22,602
2009	20,504
2010	20,512
2011	21,098
2012	35,598
2013	28,014
2014	29,715
2015	25,032
2016	20,081
2017	28,581
2018	33,243
2019	25,162
2020	49,801
2021	28,772




Approximately 70% of WMU is a CWD DMA in 2019 (Former DMA1)

**Trend** 


Stable

Year	Total
2008	0.40
2009	0.37
2010	0.43
2011	0.32
2012	0.35
2013	0.38
2014	0.34
2015	0.31
2016	0.33
2017	0.35
2018	0.28
2019	0.28
2020	0.30



FOREST HEALTH Plot - Plot Regeneration - Plot - Plot Deer Impact - Mean Deer Impact 3 or less

Year	% Adequate
2003-07	75%
2004-08	74%
2005-09	72%
2006-10	73%
2007-11	72%
2008-12	66%
2009-13	67%
2010-14	75%
2011-15	58%
2012-16	52%
2013-17	60%
2014-18	65%
2015-19	63%



Citizen Survey Results 2019 (2011) Too High 19%(5%) Just Right 53%(58%) Too Low 23%(25%)

Antlerless Allocation Options							
	Increase	Stable	Decrease				
Firearm Season Option	Harvest	Harvest	Harvest				
14 day concurrent	31,000	22,000	17,000				

<sup>&</sup>lt;sup>1</sup> - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	io characteri	Stics	
WMU	5B	% Developed	l % Forest	%Ag/Field	%Public	Area (sq mi)
		19%	28%	49%	2%	2.640

								Antle	ered H	arves	t					
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	13 000											
2005	7,381	11,717	56,000	4.6	12,000											
2006	6,995	11,384	53,000	4.5												
2007	5,974	11,143	53,000	4.6	10,000										ı	_
2008	6,762	11,184	51,000	4.4						_			- 1	ı I	ı	
2009	6,007	11,321	51,000	4.5	8,000					-				Н	Н	_
2010	6,902	12,543	50,812	4.1			_		_ =		I _			Н		
2011	7,174	12,943	50,000	3.9	6,000				Ш	┸	ш	_		Ш		
2012	8,503	12,519	51,000	4.1	.,									Н		
2013	7,443	12,847	50,000	3.9	4.000									Н		
2014	6,908	12,368	49,000	4.0	4,000											
2015	8,009	11,451	50,000	4.4										Н		
2016	8,886	12,364	50,000	4.1	2,000			-		$\blacksquare$	Н	-		Н	Н	$\dashv$
2017	8,990	12,794	57,000	4.4												
2018	9,165	14,191	58,000	4.1	- +	┻┼┻┼	▋Ţ▋				▋▃▋		_	_	Ļ	$\vdash$
2019	10,151	14,844	67,000	4.5	20	\$ 706 VQ	51 20°	200° 20	19 <sub>0</sub> 07	\$ 3	3 1/4	05 50	<sup>(</sup> ک و	2020	2029	202
2020	9,556	16,407	60,000	3.6	~	\$\psi_{\psi}\$	γ '	v v	, % <sub>-</sub>	8 V	δ, .	8 8	2	$\sim$	δ,	V

RED=7-day season

	POST-HUNT Deer Population	Trend	Stable
--	---------------------------	-------	--------

Year	Total
2008	54,020
2009	59,568
2010	53,213
2011	55,951
2012	60,723
2013	75,260
2014	63,591
2015	60,538
2016	66,282
2017	73,573
2018	85,790
2019	77,893
2020	76,623
2021	91,713




Approximately 12% of WMU is a CWD DMA in 2019 (former DMA1)

**Trend** 


Stable

Year	Total
2008	0.44
2009	0.42
2010	0.41
2011	0.40
2012	0.42
2013	0.41
2014	0.40
2015	0.37
2016	0.37
2017	0.41
2018	0.37
2019	0.33
2020	0.36



FOREST HEALTH Plot - Plot Regeneration - Plot - Plot Deer Impact - Mean Deer Impact 3 or less

Year	% Adequate
2003-07	53%
2004-08	52%
2005-09	48%
2006-10	46%
2007-11	47%
2008-12	52%
2009-13	54%
2010-14	38%
2011-15	55%
2012-16	51%
2013-17	49%
2014-18	52%
2015-19	46%



Citizen Survey Results 2019 (2011) Too High 19% (13%) Just Right 51% (58%) Too Low 20% (21%)

Antlerless Allocation Options							
Increase Stable Decrease Firearm Season Option Harvest Harvest Harvest							
7 day antlered & 7 day concurrent	72,000	62,000	51,000				
14 day concurrent	70,000	60,000	49,000				

 $<sup>^{1}</sup>$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.

			VVIV	io characteri	Stics	
WMU	5C	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		27%	37%	31%	1%	1,982

#### **Deer Harvest Antlered Harvest** Antlered Antlerless Allocation Lic/Deer<sup>1</sup> Year 12,000 2005 7,701 17,589 71,000 3.9 2006 16,123 79,000 7,708 4.8 10,000 2007 6,526 18,864 84,000 4.3 2008 8,729 20,238 92,000 4.4 8,000 7,566 23,214 4.8 2009 113,000 23,977 121,960 2010 9,400 4.7 2011 8,928 24,234 117,000 4.4 6,000 2012 7,825 23,648 111,000 4.5 2013 8,096 21,711 103,000 4.7 4,000 2014 8,035 22,152 95,000 4.3 13,551 70,000 2015\* 7,416 5.1 2,000 2016 8,328 15,643 70,000 4.4 2017 8,846 15,644 70,000 4.4 2018 7,584 16,400 70,000 4.2 , <sup>50</sup>76 2027 on to top top top top top 2019 7,646 14,364 70,000 4.8 2020 8,352 15,194 70,000 4.6 \* WMU Boundary Change


POST-HUNT Deer Population Trend

Year Total

Harvest indices (i.e., antlered harvest, antlerless lic/deer), not PASAK model, used to monitor population trend

DEER HEALTH: Fawn to Doe Ratio <sup>2</sup>	Approximately 1% of WMU is a CWD DMA in 2018	Trend	Stable

Total
0.44
0.47
0.43
0.46
0.49
0.43
0.42
0.40
0.44
0.40
0.38
0.32
0.36



FOREST HEALTH			Regeneration Assessment	
	Plot - Plot Regeneration	Plot - Plot Deer Impact	Mean Deer Impact	

Year % Adequate

#### Forest data not considered in this developed WMU

Citizen Surv	ey Results	2019 (2011)		Too High	33%(30%)	Just Right	51%(55%)	Too Low	8%(9%)
	Antlerless Allocation Options								
			Increase	Stable	Decrease				

Harvest

70,000

Harvest

61,000

Harvest

79,000

**Firearm Season Option** 

14 day concurrent

 $<sup>^{</sup>m 1}$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

			VVIV	io characteri	Sucs	
WMU	5D	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		61%	18%	11%	0%	1,327


er Harve	est				Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer <sup>1</sup>	
2005	1,460	4,166	20,000	4.5	12,000
2006	1,315	4,074	20,000	4.7	
2007	977	5,185	20,000	3.8	10,000
2008	1,343	4,533	22,000	4.7	
2009	1,130	3,911	22,000	5.2	8,000
2010	1,144	3,721	22,000	5.1	
2011	1,156	3,827	22,000	4.7	6,000
2012	1,325	3,766	19,000	4.7	
2013	1,589	4,483	18,000	4.0	4,000
2014	1,317	3,788	18,000	4.7	4,000
2015*	2,191	5,172	24,000	4.6	
2016	2,908	6,452	30,000	4.6	2,000
2017	3,327	7,526	30,000	3.9	
2018	2,631	6,001	28,000	4.6	- +
2019	2,488	6,721	29,000	4.3	\$\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
2020	2,164	6,479	29,000	4.4	

POST-HUNT Deer Population	Trend

Year Total

Harvest indices (i.e., antlered harvest, antlerless lic/deer), not PASAK model, used to monitor population trend

Total
0.41
0.49
0.48
0.49
0.44
0.41
0.48
0.42
0.42
0.40
0.46
0.33
0.44



FOREST HEALTH			Regeneration Assessment
	Plot - Plot Regeneration	Plot - Plot Deer Impact	Mean Deer Impact

Year % Adequate

#### Forest data not considered in this developed WMU

Citizen Sur	vey Results	2019 (2011)	Too High	33%(25%)	Just Right	51%(55%)	Too Low	8%(18%)
	Antlerless Allocation Options							
		Increase	Stable	Decrease				

Harvest

29,000

Harvest

23,000

Harvest

34,000

**Firearm Season Option** 

14 day concurrent

 $<sup>^{</sup>m 1}$  - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $<sup>^{\</sup>rm 2}$  - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.