

**PENNSYLVANIA GAME COMMISSION
BUREAU OF WILDLIFE MANAGEMENT
RESEARCH DIVISION
ANNUAL PROJECT REPORT**

PROJECT CODE NO.: 06110

TITLE: Survey and Statistical Support

PROJECT JOB NO.: 11101

TITLE: Game Take and Furtaker Surveys

PERIOD COVERED: 1 July 2003 to 30 June 2004

COOPERATING AGENCIES: Bureau of Automated Technology Services (BATS), Bureau of Administrative Services

WORK LOCATION(S): Harrisburg, Pennsylvania

PREPARED BY: Christopher S. Rosenberry

DATE: 23 August 2004

Abstract: A questionnaire was mailed to a random sample of purchasers of a 2003-2004 general hunting license (19,552 questionnaires mailed) to estimate number of hunters, harvest, and hunter-days of small game species during the 2003-2004 hunting season. After 3 mailings, 53.8% responded. Overall, between 2002-2003 and 2003-2004 hunting seasons, harvests and hunter numbers increased. Twenty-one year trends in harvest and hunter participation indicate a decline for nearly all small game species. A separate questionnaire was mailed to a random sample of purchasers of a furtaker license (3,206 mailed) to estimate harvest of furbearer species and trapper-days. After 2 mailings, 70.3% responded. Overall, the harvest of furbearer species and the number of hunters/trappers decreased between the 2002-2003 and 2003-2004 seasons. Twenty-one year trends for harvests of furbearers indicate harvests have declined dramatically for most species. Junior and senior combination license holders are not included in the furtaker sample, thus some furtakers are not included in survey estimates. None of the harvest estimates in this report have been adjusted to account for combination license holders. Therefore, corrected estimates from 1999-2000 and 2000-2001 in previous reports will not correspond to the estimates in this report.

OBJECTIVES

1. To estimate the number of animals harvested, number of participants, and number of days spent hunting (hunter-days) for small game species during the 2003-2004 hunting season.
2. To estimate the number of furbearers trapped or shot and number of trappers/hunters during the 2003-2004 furbearer seasons.
3. To monitor long-term trends in harvest, number of hunters and trappers, hunter-days, and harvest per 100 hunter-days.

METHODS

In March 2004, following the close of trapping and small game hunting seasons, the names and addresses of general hunting license buyers whose license

number ended in either 01 or 51, and furtakers whose license number ended in either 1 or 6, were drawn from the duplicate licenses on file in the License Division of the Bureau of Administrative Services and from the electronic file of over the counter (OTC) sales. Photocopies of the duplicates and the OTC file were used by BATS to prepare the mailing list. BATS and Bureau of Administrative Services addressed and mailed 19,552 Game Take questionnaires and 3,206 Furtaker questionnaires. In addition to the initial mailing, 2 follow-up mailings were sent to nonrespondents of the Game Take Survey and 1 follow-up mailing was sent to nonrespondents of the Furtaker Survey.

These surveys reflect major changes of information requested from hunters and trappers from pre-1990 surveys. First, information about small game and furbearer species were separated into Game Take and Furtaker Surveys, respectively. Second, the Game Take questionnaire was expanded to include more harvestable species and the number of days of hunting per species per county. Third, harvest and hunting effort on shooting preserves were requested separately for ring-necked pheasant, quail, and ducks. Fourth, estimates of coyote harvest included those shot by hunters (Game Take Survey excluding furtaker license buyers) and those trapped or shot by furtakers (Furtaker Survey). Fifth, a cover letter to encourage response was included in all mailings.

During 1990-2000, methods used to survey small game hunters and furtakers have been the same with the following exceptions. The Game Take Survey for 1992 consisted of 2/3 the usual sample size (i.e., every third 01 or 51 license was skipped) and only 2 mailings were conducted, but a telephone survey of nonrespondents was conducted to estimate nonresponse bias. Estimates using the standard estimation techniques (Shope 1985) were similar to those obtained when incorporating nonresponse bias (Diefenbach 1993). Therefore, estimates from the 1992 survey should be comparable to results from other years. In 1996 hunters were asked to report their Canada goose harvest by season (early, regular, late), and their snow goose harvest. This change was implemented to assess the effect of special goose seasons since the regular season was closed for most of the state, and to compare our estimates to those obtained by the newly implemented Migratory Bird Harvest Information Program. Since 1998 Game Take Survey turkey hunters were to report the management unit in which they hunted instead of the county. In 2000, landowner, resident senior lifetime upgrades, and resident senior lifetime renewals were included in the total licenses sold for calculating harvests and participation. This resulted in the addition of licenses to our survey population that otherwise would not have been included. The added senior licenses have existed since 1996 for lifetime renewals and 1999 for lifetime upgrades. Therefore, estimates of Game Take Surveys from 1996-1999 likely underestimate harvest by about 2-3%. Landowner licenses represent less than 0.5% of license sales and would have had minimal effect on previous survey estimates.

Since 1999, Furtaker Surveys sampled those who purchased a furtaker license but not those who purchased junior and senior combination licenses, which include furtaker privileges. As a result of this licensing change, furtaker harvest and participation estimates beginning in 1999 are biased low compared to pre-1999 estimates. To reduce this bias, a correction factor was used to adjust harvest and participation estimates in 1999 and 2000 Furtaker Surveys (Rosenberry 2000); however, this correction was discontinued in 2001 (Rosenberry 2001) and furtaker harvest estimates since 1999 are minimum estimates that do not include junior and senior combination licenses.

Beginning in 2003-04, respondents reported harvests by Wildlife Management Units (WMUs) rather than counties. In addition, the survey form was simplified by reducing the number of possible hunting areas from 4 counties to 2 WMUs. This was done because less than 5% of hunters hunted in the 3rd or 4th county on previous surveys and there are fewer WMUs (22 WMUs vs. 67 counties).

Respondents to the Game Take Survey were post-stratified on the basis of

whether or not they had purchased special licenses or stamps, to reduce the effect of nonresponse bias on estimates (see Shope 1985). In 2001, combination license holders were added to those purchasing additional stamps. Response rates for combination license holders were calculated by identifying combination license holders based on license stamps letters from the survey file and their response or nonresponse to the survey. Nonresponse bias for the Furtaker Survey was not corrected.

I estimated (by species) total harvest, number of participants, hunter-days, and harvest per 100 hunter-days based on 1,015,572 general hunting licenses sold for the Game Take Survey, and 22,454 furtaker licenses sold for the Furtaker Survey. I estimated trends over time using Pearson product-moment correlation coefficients.

In addition to the harvest and participation estimates, bobcat and fisher sightings by archery and firearms deer hunters and spring turkey hunters were added to monitor distribution and range of these species. These results are presented in furtaker and bobcat population and management reports. Eight questions were also added to assess hunter support for a number of proposals.

RESULTS

For the Game Take and Furtaker Survey respectively, 10,005 and 2,204 useable returned questionnaires were processed. The response rates, after adjusting for undeliverable questionnaires, were 53.8% for the Game Take Survey and 70.3% for the Furtaker Survey. The response rate for the Game Take survey increased slightly, and the response rate for the Furtaker survey decreased between 2002-2003 and 2003-2004.

During 2003-2004, an estimated 3,814 hunters harvested an estimated 20,678 snow geese during 16,169 hunter-days. This represents an increase in harvests, hunter participation, and hunter-days from the 2002-2003 season, when 2,310 hunters harvested 1,792 snow geese during 6,752 hunting days.

Annual Changes

Harvests of 8 of 12 small game species increased (Table 1). The number of hunters increased for 7 of 12 small game species and hunter-days increased for 9 of 12 small game species (Tables 2 and 3). Turkey harvests, hunter-days, and hunters increased for spring seasons and decreased for fall seasons.

Harvest per 100 hunter-days increased for 6 of 12 small game species (Table 4).

The number of hunters/trappers of furbearers increased or remained the same for 5 of 9 species (Table 5). Harvests decreased for 7 of 9 species (Table 6).

Twenty-one year Trends

Harvests have declined ($P < 0.10$) for nearly all species except turkey, geese, and ducks ($P < 0.10$). Trends for crow ($P = 0.09$), woodchuck ($P = 0.08$), and quail ($P = 0.25$) were not significant (Table 1). Number of hunters has declined for all seasons/species ($P < 0.10$) except spring turkey, ducks, and quail (Table 2).

Number of hunters/trappers of nearly all furbearer species has remained stable since 1990, although the number of hunter/trappers pursuing raccoons has declined ($P < 0.01$) since 1983. The number of hunters and trappers pursuing coyotes continues to increase ($P < 0.01$) (Table 5). The harvest of all furbearers for which we have 1983-2003 data has declined ($P < 0.10$). Since 1990,

coyote harvests have increased ($P < 0.01$) and weasel harvest have varied with no consistent change ($P = 0.50$).

Survey Questions

Eight questions on various topics asked respondents to record their level of support from strongly support to strongly against. Results that are weighted for purchasers and nonpurchasers are presented in Table 7. The sampling error for these questions is a maximum of plus or minus 1%.

Six percent of hunters indicated they had applied to hunt geese at Pymatuning or Middle Creek Wildlife Management Areas in the past 5 years and 12% of junior hunters indicated that they participated in the youth pheasant hunt.

RECOMMENDATIONS

1. The Game Take and Furtaker Surveys are the best source for harvest and participant data; thus, I recommend continuing these surveys.

2. Inability to sample sufficient number of combination licenses holders reduces reliability of furtaker estimates. A computerized license database would increase our ability to improve our sampling of furtakers.

3. If the response rate for the Furtaker Survey becomes <70% I recommend conducting a third mailing.

4. Following improvements to the survey instrument, the response rate for the Game Take Survey increased slightly this year. I recommend continued evaluation of the survey instrument and methodologies to improve response rate.

5. Major changes to the Game Take and Furtaker Surveys should only be instituted when more efficient sampling of license buyers, such as a computerized licensing system, is possible. If the Game Commission would implement a computerized licensing system, samples that are stratified by license type and location of residence would provide more accurate and precise harvest estimates. If such a system was implemented, phone calls to obtain estimates of harvest and hunting effort of nonrespondents to the mail survey may be useful. A computerized licensing system would allow us to survey hunters much sooner after hunting seasons ended, which has been shown to result in more accurate estimates of harvest and hunter participation (e.g., Barker 1991). Moreover, a computerized license system would provide greater flexibility in adapting sampling methods to future licensing changes that may reduce the reliability of estimates.

LITERATURE CITED

- Barker, R. J. 1991. Nonresponse bias in New Zealand waterfowl harvest surveys. *Journal of Wildlife Management* 55:126-131.
- Diefenbach, D. R. 1993. Game Take Survey. Annual Job Report. Pennsylvania Game Commission, Harrisburg, Pennsylvania, USA.
- Rosenberry, C. S. 2000. Game Take Survey. Annual Job Report. Pennsylvania Game Commission, Harrisburg, Pennsylvania, USA.
- Rosenberry, C. S. 2001. Game Take Survey. Annual Job Report. Pennsylvania Game Commission, Harrisburg, Pennsylvania, USA.
- Shope, W. K. 1985. Game Take Survey. Annual Job Report. Pennsylvania Game Commission, Harrisburg, Pennsylvania, USA.

Table 1. Harvest, by species, 1983-2003, Pennsylvania.

Year	Spring turkey	Fall turkey	Rabbits	Grouse	Squirrel	Pheasant ^a	Woodcock	Quail ^a	Dove	Geese	Ducks ^a	Hare	Woodchuck	Crow
1983	10,852	20,494	2,156,565	493,737	2,259,320		186,319		1,690,158	68,333		10,867		
1984	9,723	15,844	1,939,399	475,960	2,256,311		170,296		1,402,180	64,452		13,989		
1985	14,197	18,217	2,137,737	511,271	2,428,683		137,183		1,443,109	56,233		14,749		
1986	16,155	26,763	2,092,910	536,553	2,833,061		165,685		1,531,868	69,748		13,189		
1987	14,674	28,346	1,764,744	484,016	2,364,596		175,124		1,374,110	68,541		14,412		
1988	14,659	22,515	1,930,737	523,271	2,313,153		165,590		1,520,322	49,573		8,488		
1989	17,154	21,669	1,696,712	410,371	2,206,719		143,502		1,209,438	78,821		7,595		
1990	17,472	25,527	1,672,360	353,647	2,044,264	302,276	50,918	7,879	1,022,402	72,901	98,026	3,615	1,299,647	355,492
1991	16,606	31,979	1,462,270	293,891	1,632,108	269,065	53,183	3,005	968,421	69,127	87,478	3,579	1,304,020	257,009
1992	18,180	21,468	1,488,850	254,539	1,761,285	261,541	51,246	1,236	734,707	78,883	93,687	3,961	1,157,090	185,192
1993	24,068	30,477	1,160,939	272,690	1,585,368	250,149	52,959	4,837	735,089	84,251	133,354	2,114	1,274,166	191,639
1994	28,558	39,094	1,025,319	304,162	1,826,618	236,698	29,654	2,902	669,459	102,979	128,164	3,352	1,284,819	247,219
1995	36,401	49,748	1,010,938	315,197	1,599,104	250,930	28,624	1,204	670,791	64,382	156,511	2,997	1,225,101	295,962
1996	33,726	35,787	807,072	218,256	1,442,560	215,502	26,846	3,387	603,114	96,910	151,142	1,582	1,149,995	275,541
1997	30,956	37,398	827,520	187,770	1,352,038	219,864	23,878	1,766	506,677	115,506	188,034	1,432	1,251,145	184,944
1998	32,661	33,628	911,003	183,468	1,331,051	216,669	31,602	241	562,348	131,831	146,050	2,507	1,204,582	247,047
1999	37,806	40,718	715,862	177,355	1,236,108	211,257	25,704	3,938	519,116	128,385	164,328	2,412	1,117,970	209,273
2000	43,815	44,865	770,841	145,525	1,276,009	233,537	31,199	4,373	478,602	194,480	185,185	1,747	1,191,114	219,773
2001	49,186	48,008	701,551	159,610	1,276,603	244,282	32,504	4,276	460,971	197,767	143,907	4,584	1,187,114	195,273
2002	41,147	37,346	602,234	118,577	1,002,309	205,696	31,167	1,064	462,538	173,391	169,828	1,369	1,267,265	217,068
2003	42,876	31,100	588,310	106,587	1,063,996	234,196	42,434	2,059	500,980	228,310	191,132	1,908	1,171,888	207,707
r ^b	0.953	0.776	-0.973	-0.955	-0.937	-0.724	-0.857	-0.333	-0.944	0.861	0.835	-0.835	-0.482	-0.470
P	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.25	<0.01	<0.01	<0.01	<0.01	0.08	0.09

^aEstimates exclude harvest on shooting preserves.^bPearson product-moment correlation coefficient.

Table 2. Hunters, by species, 1983-2003, Pennsylvania.

Year	Spring turkey	Fall turkey	Rabbits	Grouse	Squirrel	Pheasant ^a	Woodcock	Quail ^a	Dove	Geese	Ducks ^a	Hare	Woodchuck	Crow
1983	255,982	367,657	738,970	471,640	614,324		148,887		188,727	70,019		28,960		
1984	209,717	322,347	626,892	419,367	525,670		120,643		162,779	66,406		27,133		
1985	214,331	298,055	619,220	423,393	528,599		100,270		150,904	62,742		25,141		
1986	246,039	336,225	612,424	442,897	552,336		110,886		166,139	65,087		27,557		
1987	206,039	282,761	516,281	374,741	472,250		96,936		137,402	50,804		19,573		
1988	226,008	300,055	528,615	390,192	472,841		93,110		143,981	53,475		21,873		
1989	224,138	296,139	497,463	365,211	464,434		87,053		131,321	43,603		17,568		
1990	191,442	234,911	436,961	299,534	369,848	274,957	30,045	5,378	93,532	33,509	28,443	7,831	123,204	39,579
1991	179,202	252,210	405,004	292,418	348,868	254,051	24,681	3,279	86,377	36,032	29,247	7,601	118,257	39,014
1992	186,738	212,104	373,800	254,724	329,726	217,189	25,916	1,444	76,998	38,301	29,263	6,156	114,515	34,442
1993	201,060	222,780	347,129	242,398	311,103	198,657	23,452	2,657	73,462	41,577	35,782	5,801	109,576	34,648
1994	224,405	244,095	335,715	259,727	326,271	205,384	19,401	1,323	74,589	40,106	34,097	7,236	117,251	37,841
1995	239,521	261,395	297,570	239,014	293,852	182,224	15,702	1,451	67,754	28,715	30,274	5,949	113,127	36,782
1996	241,613	250,377	280,351	214,272	279,259	171,275	14,464	1,184	65,808	31,119	32,434	5,011	101,576	30,087
1997	233,287	249,934	261,115	197,994	267,051	148,900	13,374	1,009	60,178	30,574	32,180	3,723	104,561	30,696
1998	194,819 ^b	199,696 ^b	242,509	183,511	252,738	158,497	12,907	1,116	57,579	32,871	34,103	5,506	92,517	31,390
1999	237,984	244,638	221,179	174,576	238,887	142,142	12,212	1,550	49,551	33,734	31,503	4,379	90,853	29,131
2000	231,860	230,448	229,906	162,073	238,540	149,260	12,977	1,870	52,496	35,628	31,998	3,666	99,294	29,371
2001	230,115	228,564	213,295	161,186	231,436	146,751	14,411	2,029	51,144	38,292	31,893	4,930	99,787	33,343
2002	218,931	217,099	195,078	149,106	201,694	123,879	12,652	1,342	50,883	41,240	32,328	3,818	91,149	28,470
2003	246,820	211,967	181,426	134,115	199,922	130,676	15,321	3,518	46,580	44,467	34,173	5,091	92,986	27,591
r ^c	0.150	-0.823	-0.976	-0.978	-0.966	-0.940	-0.877	-0.341	-0.940	-0.733	0.448	-0.880	-0.901	-0.851
P	0.52	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.23	<0.01	<0.01	0.11	<0.01	<0.01	<0.01

^aEstimates exclude number of hunters on shooting preserves.

^bCautionary note: these low values may have been caused by inadvertently not including the TMA map on the 1998-1999 survey instructions. See 1998-1999 annual report.

^cPearson product-moment correlation coefficient.

Table 3. Hunter-days , by species, 1990-2003, Pennsylvania.

Year	Spring turkey	Fall turkey	Rabbits	Grouse	Squirrel	Pheasant ^a	Woodcock	Quail ^a	Dove	Geese	Ducks ^a	Hare	Woodchuck	Crow
1990	861,086	872,815	2,901,567	1,764,129	2,345,050	1,287,702	133,947	24,493	475,402	171,436	141,411	15,632	1,228,548	223,525
1991	781,499	851,155	2,474,017	1,580,574	2,004,826	1,115,902	119,238	13,630	409,149	167,342	132,775	15,397	1,341,605	227,527
1992	799,621	696,705	2,210,784	1,331,444	1,814,807	902,308	97,699	3,228	329,087	188,303	135,656	11,650	1,191,725	170,185
1993	843,987	753,896	1,926,331	1,246,856	1,721,261	859,018	94,588	16,683	326,265	202,644	174,023	11,882	1,338,167	201,412
1994	1,003,939	857,959	2,104,454	1,438,808	1,919,013	937,974	73,958	4,455	340,661	217,021	163,690	15,208	1,294,150	209,854
1995	1,084,725	865,565	1,769,363	1,281,923	1,630,631	844,056	62,819	6,022	295,114	128,611	165,196	11,712	1,253,239	193,952
1996	1,103,556	867,072	1,641,774	1,130,129	1,568,102	733,806	51,493	5,061	280,603	165,523	168,834	9,230	1,246,439	186,781
1997	1,019,546	834,253	1,525,740	1,022,603	1,462,230	648,985	48,577	2,837	237,910	214,269	199,017	6,849	1,241,112	178,724
1998	881,026 ^b	691,787 ^b	1,517,673	994,150	1,422,957	775,398	55,343	6,704	261,442	212,538	188,694	11,805	1,359,595	222,980
1999	1,023,988	807,292	1,268,639	882,167	1,306,098	605,034	47,142	5,004	207,743	230,635	189,306	6,864	1,151,067	173,186
2000	995,472	780,297	1,295,397	817,545	1,254,598	652,602	56,098	8,906	230,991	259,153	202,279	5,351	1,196,679	157,828
2001	1,025,011	800,113	1,319,445	894,983	1,371,514	714,970	66,333	8,355	217,529	284,517	183,880	10,837	1,280,855	250,869
2002	964,575	770,899	1,043,657	723,845	1,069,972	520,372	52,222	9,638	209,960	277,528	210,663	8,761	1,178,530	164,521
2003	1,069,299	757,304	1,058,453	700,729	1,049,995	595,908	75,627	13,834	210,869	331,784	226,495	11,206	1,103,755	237,168
r ^c	0.617	-0.311	-0.957	-0.961	-0.954	-0.893	-0.738	-0.284	-0.918	0.823	0.919	-0.648	-0.490	-0.062
P	0.02	0.28	<0.01	<0.01	<0.01	<0.01	<0.01	0.32	<0.01	<0.01	<0.01	0.01	0.08	0.83

^aEstimates exclude effort on shooting preserves.

^bCautionary note: these low values may have been caused by inadvertently not including the TMA map on the 1998-1999 survey instructions. See 1998-1999 annual report.

^cPearson product-moment correlation coefficient.

Table 4. Harvest per 100 hunter-days, by species, 1990-2003, Pennsylvania.

Year	Spring turkey	Fall turkey	Rabbits	Grouse	Squirrel	Pheasant ^a	Woodcock	Quail ^a	Dove	Geese	Ducks ^a	Hare	Woodchuck	Crow
1990	2.0	2.9	57.6	20.0	87.2	23.5	38.0	32.2	215.1	42.5	69.3	23.1	105.8	159.0
1991	2.1	3.8	59.1	18.6	81.4	24.1	44.6	22.0	236.7	41.3	65.9	23.2	97.2	113.0
1992	2.3	3.1	67.3	19.1	97.1	29.0	52.5	38.3	223.3	41.9	69.1	34.0	97.1	108.8
1993	2.9	4.0	60.3	21.9	92.1	29.1	56.0	29.0	225.3	41.6	76.6	17.8	95.2	95.1
1994	2.8	4.6	48.7	21.1	85.2	25.2	40.1	65.1	196.5	47.5	78.3	22.0	99.3	117.8
1995	3.4	5.7	57.1	24.6	98.1	29.7	45.6	20.0	227.3	50.1	96.8	25.6	97.8	152.6
1996	3.1	4.1	49.2	19.3	92.0	29.4	52.1	66.9	214.9	55.3	89.5	17.1	92.3	147.5
1997	3.0	4.5	54.2	18.4	92.5	33.9	49.2	62.2	213.0	53.9	94.5	20.9	100.8	103.5
1998	3.7	4.9	60.0	18.5	93.5	27.9	57.1	3.6	215.1	66.9	77.4	21.2	88.6	110.8
1999	3.7	5.0	56.4	20.1	94.6	34.9	54.5	78.7	249.9	55.7	86.8	35.1	97.1	120.8
2000	4.4	5.7	59.5	17.8	101.7	35.8	55.6	49.1	207.2	75.0	91.5	32.6	99.5	139.2
2001	4.8	6.0	53.2	17.8	93.1	34.2	49.0	51.2	211.9	67.2	78.3	42.3	92.7	77.8
2002	4.3	4.8	57.7	16.4	93.7	39.5	59.7	11.0	220.3	62.5	80.6	15.6	107.5	131.9
2003	4.0	4.1	55.6	15.2	101.3	39.3	56.1	14.9	237.6	68.8	84.4	17.0	106.2	87.6
r ^b	0.930	0.639	-0.236	-0.596	0.586	0.906	0.657	-0.029	0.074	0.909	0.528	0.081	0.133	-0.317
P	<0.01	0.01	0.41	0.02	0.3	<0.01	0.01	0.92	0.80	<0.01	0.05	0.78	0.65	0.27

^aEstimates exclude effort on shooting preserves.

^bPearson product-moment correlation coefficient.

Table 5. Furbearer hunters and trappers, by species, 1990-2003, Pennsylvania.

Year	Raccoon	Muskrat	Red Fox	Gray fox	Opossum	Skunk	Mink	Coyote ^a	Weasel
1990	9,676	4,147	7,941	6,542	3,653	1,914	2,560	7,782	508
1991	9,921	4,865	7,827	6,613	3,915	2,264	2,726	12,184	422
1992	9,525	4,419	7,019	6,263	3,793	2,208	2,539	13,643	452
1993	8,195	4,227	6,790	6,089	3,369	1,967	2,465	14,260	387
1994	7,066	5,570	8,319	7,515	4,267	3,071	3,212	20,597	784
1995	9,718	4,465	8,080	6,908	3,989	2,643	2,879	20,413	853
1996	12,951	6,478	10,007	8,361	6,140	3,443	3,703	21,937	942
1997	13,750	7,363	10,330	8,553	6,386	3,473	4,434	24,526	1,125
1998	12,794	5,900	9,982	8,594	5,558	2,948	3,512	30,016	733
1999 ^b	7,555	3,230	6,996	6,061	2,653	1,718	2,152	28,265	392
2000 ^b	6,996	3,121	7,280	6,353	2,870	1,750	2,026	28,270	509
2001 ^b	7,935	3,997	8,234	6,938	3,180	2,036	2,587	36,249	619
2002 ^b	7,295	3,287	8,022	6,494	3,434	2,116	2,433	28,535	676
2003 ^b	7,292	3,362	6,998	5,547	3,585	2,132	2,305	29,048	453
r ^c	-0.778 ^d	-0.323	0.028	-0.079	-0.145	-0.117	-0.162	0.928	0.092
P	<0.01	0.26	0.92	0.79	0.62	0.69	0.58	<0.01	0.75

^aCombines estimates from Game Take Survey and Furtaker Survey.

^bCautionary note: Estimates are minimum estimates that do not account for combination licenses.

^cPearson product-moment correlation coefficient.

^dCorrelation coefficient estimated using 1983-2003 data.

Table 6. Furbearer harvests, by species, 1983-2003, Pennsylvania.

Year	Raccoon	Muskrat	Red fox	Gray fox	Opossum	Skunk	Mink	Coyote ^{a,b}	Weasel ^a
1983	449,499	575,530	88,643	64,754	339,436	86,769	13,089		
1984	495,106	621,111	75,532	66,975	339,294	72,050	23,627		
1985	557,989	362,074	68,074	40,476	237,493	48,847	13,932		
1986	426,625	440,880	95,330	46,387	210,953	39,064	16,008		
1987	443,934	346,558	74,590	56,944	217,552	39,632	18,513		
1988	247,743	230,058	52,778	23,102	105,881	16,371	12,914		
1989	155,761	141,577	43,525	28,818	80,660	20,409	9,669		
1990	116,443	112,358	32,699	21,653	36,574	9,298	7,053	1,810	798
1991	130,608	156,014	28,495	30,409	37,177	8,907	10,355	3,719	481
1992	124,404	135,533	27,611	25,395	27,754	7,221	9,157	4,402	343
1993	118,964	121,657	25,862	23,839	25,807	7,920	7,808	6,161	526
1994	186,551	178,145	30,649	34,691	29,621	12,620	10,208	6,240	723
1995	120,462	130,442	31,110	23,518	29,688	9,995	8,602	6,662	687
1996	214,958	146,013	29,623	23,307	48,549	11,571	9,315	7,957	589
1997	194,696	216,066	36,923	26,043	60,717	12,344	14,063	6,685	1,172
1998	195,110	148,202	47,202	32,922	56,287	11,190	12,238	11,652	662
1999 ^d	96,270	88,426	34,297	21,762	28,950	6,853	12,512	8,797	336
2000 ^d	97,509	79,933	30,893	20,096	25,062	7,248	7,980	10,160	313
2001 ^d	121,810	121,994	33,003	23,275	27,192	9,245	13,214	12,363	815
2002 ^d	106,485	75,340	33,007	18,805	34,787	7,207	10,069	11,444	406
2003 ^d	104,781	71,368	31,592	15,956	33,760	9,319	6,494	11,697	359
r ^c	-0.778	-0.798	-0.736	-0.765	-0.787	-0.758	-0.527	0.943	-0.197
P	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	0.01	<0.01	0.50

^aNo data are available prior to 1990.

^bCombines estimates from the Game Take and Furtaker surveys.

^cPearson product-moment correlation coefficient.

^dCautionary note: Estimates are minimum estimates that do not account for combination licenses.

Table 7. Responses to opinion questions, 2003-2004 Game Take Survey.

Question	Strongly Support	Support	Undecided	Against	Strongly Against
1. Either sex pheasant hunting on all State Game Lands.	22%	28%	26%	14%	9%
2. Expanding either sex pheasant hunting in areas with marginal habitat to increase hunting opportunities of Commission produced pheasants.	20%	30%	30%	13%	8%
3. September goose hunting at Pymatuning and Middle Creek Wildlife Management Areas.	20%	28%	47%	2%	2%
4. August seasons to control Canada goose populations.	32%	33%	29%	4%	2%
5. Spring seasons to control Canada and greater snow goose populations.	30%	32%	30%	5%	3%
6. Hunting ½ hour after sunset to control Canada and greater snow goose populations.	25%	25%	33%	11%	6%