

**PENNSYLVANIA GAME COMMISSION  
BUREAU OF WILDLIFE MANAGEMENT  
PROJECT ANNUAL JOB REPORT**

**PROJECT CODE NO.:** 06210

**TITLE:** White-tailed Deer Research/Management

**JOB CODE NO.:** 21001

**TITLE:** Deer Health, Forest Habitat Health, Deer Harvests, and Deer Population Trends by Wildlife Management Unit

**PERIOD COVERED:** 1 July 2009 through 30 June 2010

**COOPERATING AGENCIES:** Pennsylvania Cooperative Fish and Wildlife Research Unit, Pennsylvania Department of Conservation and Natural Resources, Pennsylvania State University, and U.S. Forest Service

**WORK LOCATION(S):** Statewide

**PREPARED BY:** Christopher Rosenberry, Bret Wallingford, and Jeannine Tardiff Fleegle

**DATE:** 1 June 2010

**ABSTRACT** We monitored Wildlife Management Unit (WMU) deer health, forest habitat health, and deer population trends using reproductive parameters from road-killed does, advanced tree seedling and sapling regeneration from the Pennsylvania Regeneration Study, deer harvest estimates and compositions, and field studies. Deer health was judged to be “above target” in 6 WMUs, “below target” in 1 WMU, and “at target” in 15 WMUs. Forest habitat health was judged to be good in 1 WMU, fair in 19 WMUs, and poor in 1 WMU. Hunters harvested 308,920 deer (108,330 antlered and 200,590 antlerless) in the 2009-10 deer seasons. Deer populations in most WMUs remained stable. Seasons and antlerless allocations recommendations to stabilize deer populations in nearly all WMUs were modified by the Board of Commissioners to increase deer populations in 20 of 22 WMUs.

## **OBJECTIVE**

To monitor deer health, forest habitat health, deer harvests, and deer population trends by Wildlife Management Unit (WMU).

## **METHODS**

### **Deer Health**

To obtain data on deer health, Wildlife Conservation Officers (WCOs) and other personnel examined female deer killed by various causes from 1 February through 31 May 2009. They recorded location (county, township, and WMU), date killed, cause of death, and number and sex of embryos for each doe on a form attached to a deer jaw envelope. One side of the lower jaw was removed from each deer for age determination. Jaws were forwarded to Regional Wildlife Management Supervisors, who forwarded them to the Deer and Elk Section for ageing in June 2009. Personnel in the Bureau of Automated Technology Services (BATS) processed the reproductive data and provided summary reports for the state and each WMU.

Based on results from published studies (Cheatum and Severinghaus 1950, Verme 1965, Verme 1967, Verme 1969, Hesselton and Sauer 1973, Hesselton and Jackson 1974, McCullough 1979, Stoll and Parker 1986, Folk and Klinstra 1991, Osborne et al 1992, Taylor 1996, Swihart et al 1998), we used embryo counts per adult female (2-years-old and older) to assess deer health in each WMU. We chose a target of 1.50 embryos per adult female for deer to be considered healthy. Because of difficulties in gathering sufficient samples for 1-year-old females, 2-year-old females, and 3-year-old and older females, we combined 2-year-old and 3-year-old and older groups into a single group (“adult female”). This grouping also makes our data set comparable to previously published work. The value of 1.50 was chosen for the following reasons: 1) 1.50 embryos per adult female (2-year-old and older female) corresponds to a population producing at a high and sustainable level (Downing and Guynn 1985), 2) based on more than a dozen studies from the United States and Canada, a value of 1.50 embryos per adult female represents the middle ground between deer with low and high nutrition, and 3) 1.50 embryos per adult doe is achievable for WMUs in Pennsylvania. Studies used to assess the suitability of 1.50 embryos per adult female come from states and provinces including Michigan, Manitoba, Ohio, New York, and Pennsylvania.

Because our reproduction point estimate is based on a sample, we conducted a two-tailed  $t$  test for differences between the sample mean and our target of 1.50 embryos per adult female. We believe this approach is better than using a strict cutoff because it provides a framework to assess not only the point estimates, but also variation.

*Decision Rules Used to Determine Deer Health.--*

1. Does 3-year estimate of embryos per adult female have a coefficient of variation (CV) of  $\leq 13\%$ <sup>a</sup>?

a. YES. Is the WMU’s point estimate of embryos per adult female statistically different from 1.50?

i. NO. Deer health is “at target”.

ii. YES. Is the point estimate above 1.50?

1. YES. Deer health is “above target”.

2. NO. Deer health is “below target”.

b. NO. Larger sample sizes are needed to achieve desired levels of estimator precision. Deer health will be assessed based on point estimate and small sample size noted.

<sup>a</sup> – A coefficient of variation (CV) of approximately <13% is considered sufficient for accurate population management (Skalski and Millsbaugh 2002, Skalski et al. 2005, Millsbaugh et al. 2006). At this time, it typically requires pooling of 3 years of data to achieve CVs of less than 13%.

### **Forest Habitat Health**

We used forest regeneration to assess forest habitat health. Forest regeneration is not just a measure for the benefit of the forest, but also for deer and wildlife. For deer, seedling and sapling trees provide food and cover. As a result, measuring regeneration is an important measure of the sustainability of a forest, and available food and cover that benefit deer and other wildlife.

To obtain data on forest regeneration, advanced tree seedling and sapling regeneration (ATSSR) data are collected as part of a systematic sampling scheme from public and private lands in WMUs from the Pennsylvania Regeneration Study. This study is being conducted as part of the Forest Inventory Analysis (FIA) by Pennsylvania Department of Conservation and Natural Resources (DCNR), Pennsylvania State University (PSU), and U.S. Forest Service (USFS). Subsets of all plots are collected each year, with a complete sampling of plots occurring every 5 years. ATSSR from 2 groupings of tree species are available from the Pennsylvania Regeneration Study. The measure selected for use in deer management is the grouping of dominant canopy species and species capable of achieving high canopy status. “The composition of the ATSSR has a direct impact on the future composition of the forest overstory (Marquis and others 1994). To cover the range of future forest character and client needs 2 composition groupings are used. The first groups tree species by preference for timber management. The second composition grouping represents the forest’s ability to regenerate the existing dominant canopy. Dominant species include those that contribute at least 2% of the State’s total-tree biomass and are able to grow into the existing canopy; Other High Canopy species include all others that are capable of attaining canopy dominance” (McWilliams et al. 2004).

We requested ATSSR data for dominant canopy species and species capable of achieving high canopy status by WMU from the USFS and DCNR. Determination of adequate regeneration was based on levels of deer browse impact observed in the area of each plot. For example, a greater number of saplings are required to replace the existing canopy where deer impact is “very high” compared to fewer saplings required where deer impact is “very low”. The scaled levels of deer impact indicate deer population size in relation to food availability in a given area (i.e., carrying capacity). Areas with ample food to support the local deer population will be evident by very low to medium deer impact. Areas lacking food to support the local deer population will be evident by high to very high deer impact. These critical stocking guidelines were derived from extensive literature reviews and decades of research on deer-habitat interactions (Marquis et al. 1992). In 2008 we began using browse impact and associated stocking levels in the habitat health measure. Because of the sampling scheme used in the Pennsylvania Regeneration Study, it takes 5 years to visit all sample plots.

Based on input from cooperating agencies that designed and conduct the Pennsylvania Regeneration Study and an internal Game Commission review of the forest habitat health measure, we defined forest habitat as “good” if 70% or more of the sampled plots contained adequate regeneration. If less than 50% of the plots contained adequate regeneration, forest habitat health was considered “poor”. “Fair” falls between levels for “good” and “poor”.

Similar to the deer health measure, the forest habitat health measure is based on a sample of plots from across a WMU and we use a statistical test to assess regeneration levels. By using a statistical test to assess differences from predetermined levels (e.g., 70%), we take into account both the point estimate and associated variation.

When data are collected according to proper sampling design, estimates can be statistically compared to 50% and 70% levels using a t-test. The t-test determines whether the estimate is different from the 50% or 70% level based on standard statistical procedures. Since reliability of statistical tests is related to sample sizes, forest habitat health determinations are made based on 5-year data sets to maximize sample size and reliability of statistical tests.

*Decision Rules Used to Determine Forest Habitat Health.*--We developed a set of criteria to assign a value of “good”, “fair”, or “poor” for forest habitat health. A WMU’s forest habitat health was considered “good” if the observed percentage of plots with adequate regeneration was greater than, equal to, or not significantly different than 70%. If a WMU’s forest habitat health was not significantly different from 70% and not significantly different from 50%, then forest habitat health was considered “fair”. A WMU’s forest habitat health also was considered “fair” if: 1) the observed percentage of plots with adequate regeneration was equal to 50%; or 2) between 50% and 70% and significantly less than 70%; or 3) not significantly different than 50%. A WMU’s forest habitat health was considered “poor” if the observed percentage of plots with adequate regeneration was significantly less than 50%.

### **Deer Harvest Estimates and Composition**

To estimate deer harvests and collect data for monitoring deer population trends, 33 data collection teams examined deer in assigned areas across the state. Each team collected data for 3 days during the first week of the regular firearms season, 2 days during the second week of the season, and 2 days after the close of the season. Data were recorded electronically on Pendragon Forms 5.1 software using a Windows Mobile hand-held computer (Trimble Nomad), and downloaded to a Harrisburg data collection point. Data collected included age, sex, location of harvest (WMU, county, and township), and hunting license number from ear tags. Deer teams determined deer age as 6 months (fawn), 18 months (yearling), or at least 30 months (adult) using tooth wear and replacement (Severinghaus 1949). Data collection teams also recorded points of antlers to determine antler characteristics by age class.

A data entry company was contracted to enter deer harvest report card data. BATS validated and processed harvest data and ran harvest data analysis programs. For each WMU the analyses included: the number of antlered and antlerless deer checked by aging teams, the number of antlered and antlerless deer checked by deer aging teams and reported by hunters, the total number of antlered and antlerless deer reported by hunters, age and sex composition of the harvest, and reported regular firearms, muzzleloader, and archery harvests.

Deer harvests were estimated using mark-recapture methods. When estimating deer harvests, we used a closed, 2-sample Lincoln-Petersen estimator where deer were considered marked when they were checked in the field by deer aging teams. Recapture occurred when marked deer were reported on report cards sent in by hunters.

Because reporting rates in Pennsylvania vary by year, antlered and antlerless deer, and WMU (Rosenberry et al. 2004), deer harvest estimates were calculated for antlered and antlerless deer in each WMU using Chapman's (1951) modified Lincoln-Petersen estimator. This estimator is recommended (Nichols and Dickman 1996) because it has less bias than the original Lincoln-Petersen estimator (Chapman 1951).

### **Deer Population Trends**

We used multiple methods to monitor deer population trends including a modified sex-age-kill (SAK) model (Eberhardt 1960, Creed et al. 1984, Skalski and Millspaugh 2002), antlerless hunter success index (i.e., estimated antlerless harvest divided by the number of antlerless licenses), and an antlered harvest index (i.e., estimated antlered harvest for a WMU).

We modified the standard SAK model to account for Pennsylvania's antler restrictions to monitor deer population trends. Modifications involve estimation of 1.5-year-old and 2.5-year-old and older male populations. Population trend monitoring relies on research data from Pennsylvania (e.g., Long et al. 2005), harvest estimates, and deer aging data. Population monitoring began with adult males (males 1.5 years of age and older) and progressed to females and fawns.

The modified SAK procedure began by estimating males 2.5 years of age and older from harvest estimates and adult male harvest rates. Once the population of males 2.5 years of age and older were estimated, we determined the 1.5-year-old male population. Because protection levels of 1.5-year-old males varied among WMUs and harvest rates could also vary, we worked back in time to generate harvest rates for 1.5-year-old males. First, we determined the pre-hunt population of 1.5-year-old males in the preceding year using current year population estimate of 2.5-year-old males, survival rate from 1.5 to 2.5 years of age, and estimated harvest of 1.5-year-old males in the preceding year. Harvest rate of 1.5-year-old males from the preceding year was then calculated using the pre-hunt population and estimated harvest of 1.5-year-old males. Current year's population of 1.5-year-old males was determined using a 3-year running average of harvest rates of 1.5-year-old males from the 3 previous years. Following determination of the 1.5-year-old males and males 2.5 years of age and older, calculation of female, fawn, and the total populations followed procedures similar to Skalski and Millspaugh (2002).

When interpreting results from the modified SAK procedure, it is important to know that due to the nature of population reconstruction methods, such as those used in the SAK procedure, the most accurate population estimate for a particular year occurs at some point in the future when data for each cohort of deer is complete (Skalski et al. 2005). Consequently, for the most recent years, population numbers should be viewed as indices rather than estimates (Skalski et al. 2005). Second, due to necessary assumptions of this population monitoring procedure, population numbers used to assess trends should be viewed as relative (i.e., whether trends are

increasing, decreasing, or remaining stable), not absolute numbers. As we accumulate more years of data and results from on-going internal and external evaluations, refinements to this procedure will occur.

Population trends are reported as changes from year to year ( $\lambda$ ) and are calculated as  $\lambda = \frac{\hat{N}_{t+1}}{\hat{N}_t}$  where  $\hat{N}_{t+1}$  is the deer population in year  $t+1$  and  $\hat{N}_t$  is the deer population in year  $t$  (Skalski et al. 2005). A value of  $\lambda = 1.00$  would indicate no change in deer population. Values greater than 1.00 indicate increases and values less than 1.00 indicate decreases. Deer management objectives and recommendations are based on population trends. As a result, we do not make management recommendations in response to individual  $\lambda$ s, but rather we based management recommendations on multi-year trends.

We identified population trends as increasing, decreasing, or stable based on graphical and statistical methods, specifically the Mann-Kendall Test for Trend (Mann 1945, Kendall and Gibbons 1990). We chose this test because it provides a statistical test of trend in data without complex calculations and does not require actual differences between years. Since effective state agency deer programs must consider public involvement and perceptions, it is important that we assess trends with a test that is statistically appropriate, utilizes information available to the public (e.g., a graph of estimates over time), and is relatively easy to explain.

## RESULTS

### Deer Health

Pennsylvania Game Commission (PGC) personnel examined 1,307 females during the 2009 pre-fawning season. Seven hundred and sixty-six were pregnant. Twenty percent of the fawns, and 89% of the adults were pregnant or lactating. Pregnant fawns averaged 1.18 embryos/female. Pregnant adults averaged 1.81 embryos/female. The average reproductive rates for pregnant and barren fawns and adults were 0.23 and 1.60 embryos/female, respectively. The average reproductive rate for all females was 1.01 embryos/doe (Table 1).

WMU deer health assessments were based on reproduction from 3 consecutive years, 2007 to 2009. We pooled these 3 years because annual sample sizes are too small to make reliable inferences. Of 22 WMUS, we identified 6 with “above target” deer health, 1 with “below target” deer health, and the remaining 15 with “at target” deer health (Table 2).

### Forest Habitat Health

WMU forest habitat health assessments were based on the 5 years of the Pennsylvania Regeneration Study from 2005 to 2009. We identified 1 WMU with good forest habitat health, 19 with fair forest habitat health, and 1 with poor forest habitat health (Table 3). One unit, 5D, does not have data to make an assessment. Results from this report can be compared to reports since the 2007-08 report, but not to previous reports because of the addition of deer browse impact in assessing regeneration adequacy.

### Deer Harvest Estimates and Composition

PGC personnel checked an average of 361 (range: 22 to 584) antlered deer and 768 (range: 163 to 1,693) antlerless deer per WMU during the 2009 firearms season (Table 4). Based on deer checked and report cards sent in by successful hunters, hunters harvested an estimated 308,920 deer in the 2009-10 deer seasons (Table 4). The antlered harvest was 108,330, a decrease of 12% from the 2008-09 harvest of 122,410. The antlerless harvest was 200,590, a decrease of 6% compared to the harvest of 213,440 in 2008-09.

Antlered harvests were composed of 49% 1.5-year-old males and 51% 2.5-year-old and older males (Table 5). Compared to years prior to implementation of antler restrictions during the 2002-03 hunting seasons, the age structure of the antlered harvest has increased, as has the number of 2.5-year-old and older bucks harvested (Table 5). Antlerless harvest composition has changed little since 1997-98 hunting seasons (Table 6).

### **Deer Population Trends**

Population changes ( $\lambda$ s) for 7 WMUs were between 0.90 and 1.10 from 2008 to 2009 based on preliminary population estimates for 2009 (Table 7). Since 2005, most WMU populations have remained relatively stable.

### **Deer Management Recommendations**

We recommended that all deer hunting seasons and regulations remain in place for the 2010-2011 hunting seasons. These regulations include a 12-day concurrent antlered and antlerless firearms season for all hunters (excluding WMUs 2D, 2G, 3C, and 4B); a 7-day antlerless muzzleloader season in October; a 3-day antlerless rifle season in October for junior, senior, disabled, and military license holders; sale of unsold antlerless licenses, up to 2 per hunter that remain after all hunters have had an opportunity to purchase one; and field possession regulations that allow a hunter to harvest another deer after tagging the first deer harvested. We also recommended a 5 day antlered and 7 day concurrent firearms season in WMUs 2D, 2G, 3C, and 4B remain in place as 2 years of data collection in a 4-year research program to evaluate the social and biological impacts of this season have been completed. We recommended that antlerless allocations remain the same in all WMUs except 3B (reduction of 8,000 licenses) and 5C (increase of 9,000 licenses). These allocations were designed to stabilize populations in most WMUs, increase the population in 3B, and decrease the population in 2B, 5C, and 5D.

### **Action by the Board of Commissioners**

The Board of Commissioners (BOC) modified the firearms season in WMUs 2C, 2E, 4D, and 4E to a 5-day antlered/7-day concurrent season split format. The BOC also modified the recommended antlerless allocations. Initial modifications included reductions in 1B (5%), 2C (9%), 2D (10%), 2F (10%), 3D (10%), 4D (20%), and 4E (10%). Further action taken by the BOC reduced all WMU antlerless allocations by the number of deer management assistance program (DMAP) permits sold in the previous year. A cap for DMAP permits was also passed whereby DMAP permits cannot exceed those sold in the previous year for each WMU. The end result was reduction of the recommended antlerless allocation in 21 WMUs, and the elimination of DMAP in 2 WMUs. The most significant reductions were in 1B (7%), 2C (10%), 2D (10%), 2F (21%), 2G (42%), 3B (21%), 3D (15%), 4A (5%), 4D (25%), and 4E (10%).

## **RECOMMENDATIONS**

1. Investigate alternative deer herd health options, including fawn:doe ratios in harvest data
2. Identify and develop additional analyses and measurements to improve the forest habitat health measure's ability to account for factors other than deer that affect forest regeneration and to most directly monitor deer impacts on forest regeneration.
3. Maintain deer aging sampling effort. Current numbers of deer checked in the field provide reasonably precise harvest estimates in most WMUs. Harvest estimates are least precise in smaller WMUs where it is more difficult to collect sufficient data.
4. Continue to evaluate validity of assumptions and population monitoring procedures through internal and external peer-review. Prioritize research needs based on internal and external reviews.
5. Continue the 5-day antlered/7-day concurrent firearms season as currently approved in WMUs 3C and 4B to allow for further collection of research data. Some objectives of the research in WMUs 2D and 2G have been severely compromised due to the BOC actions involving changes in antlerless allocations. However, research should continue in these areas to measure the effects of changes on the deer population. For research control WMUs 1A, 2F, 3A, and 4A, changes in the antlerless allocations have increased complexity and decreased validity of research analyses.
6. Return concurrent antlered and antlerless firearms seasons for all WMUs except 2D, 2G, 3C, and 4B in 2011-12 hunting seasons or increase antlerless allocations as needed. This provides more hunting opportunities to hunters and maintains consistency in hunting seasons that is important to monitoring population trends. In addition, the antlerless allocation can control the antlerless harvest without changing season length.
7. Continue antler restriction regulations in accordance with goals and objectives of the 2009-2018 deer management plan.
8. Continue to allow hunters to purchase and use the entire antlerless allocation.
9. Return to antlerless license allocations designed to achieve deer management goals as defined in the deer management plan.

## **LITERATURE CITED**

- Chapman, D. G. 1951. Some properties of the hypergeometric distribution with applications to zoological censuses. University of California Publications on Statistics 1:131-160.
- Cheatum, E. L., and C. W. Severinghaus. 1950. Variations in fertility of white-tailed deer related to range conditions. Transactions of the North American Wildlife Conference 15:170-190.

- Creed, W. A., F. Haberland, B. E. Kohn, and K. R. McCaffery. 1984. Harvest management. Pages 243-260 *in* L. K. Halls, editor. White-tailed deer ecology and management. Stackpole Books, Harrisburg, Pennsylvania, USA.
- Downing, R. L., and D. C. Guynn, Jr. 1985. A generalized sustained yield table for white-tailed deer. Pages 95-103 *in* S. L. Beasom, and S. F. Roberson, editors. Game harvest management. Caesar Kleberg Wildlife Research Institute, Kingsville, Texas, USA.
- Eberhardt, L. L. 1960. Estimation of vital characteristics of Michigan deer herds. Michigan Department of Conservation, Game Division Report 2282. East Lansing, USA.
- Folk, M. J., and W. D. Klinstra. 1991. Reproductive performance of female key deer. *Journal of Wildlife Management* 55:386-390.
- Hesselton, W. T., and L. W. Jackson. 1974. Reproductive rates of white-tailed deer in New York. *New York Fish and Game Journal* 21:135-152.
- Hesselton, W. T., and P. R. Sauer. 1973. Comparative physical condition of four deer herds in New York according to several indices. *New York Fish and Game Journal* 20:77-107.
- Kendall, M. G., and J. D. Gibbons. 1990. Rank Correlation Methods. Fifth edition. Edward Arnold, London, United Kingdom.
- Long, E. S., D. R. Diefenbach, C. S. Rosenberry, B. D. Wallingford, and M. D. Grund. 2005. Landscape structure influences dispersal distances of a habitat generalist, the white-tailed deer. *Journal of Mammalogy* 86:623-629.
- Mann, H. B. 1945. Non-parametric tests against trend. *Econometrica* 13:245-259.
- Marquis, D. A., R. L. Ernst, and S. L. Stout. 1992. Prescribing silvicultural treatments in hardwood stands of the Alleghenies. Revised editor. U.S. Forest Service General Technical Report NE-96.
- McCullough, D. R. 1979. The George Reserve deer herd: population ecology of a K-selected species. University of Michigan Press. Ann Arbor, USA.
- McWilliams, W. H., C. A. Alerich, D. A. Devlin, A. J. Lister, T. W. Lister, S. L. Sterner, and J. A. Westfall. 2004. Annual inventory report for Pennsylvania's forests: results from the first three years. Resource Bulletin NE-159. USDA Forest Service, Newtown Square, Pennsylvania, USA.
- Millsbaugh, J. J., M. S. Boyce, D. R. Diefenbach, L. P. Hansen, K. Kammermeyer, and J. R. Skalski. 2006. An evaluation of the SAK model as applied in Wisconsin. A report to the Wisconsin Department of Natural Resources. Madison, USA.

- Nichols, J. D. and C. R. Dickman. 1996. Capture-recapture methods in measuring and monitoring biological diversity: standard methods for mammals. Pages 217-226 in D. E. Wilson, F. R. Cole, J. D. Nichols, R. Rudran, and M. S. Foster, editors. Smithsonian Institution Press, Washington D.C., USA.
- Osborne, J. S., A. S. Johnson, P. E. Hale, R. L. Marchinton, C. V. Vansant, and J. M. Wentworth. 1992. Population ecology of the Blackbeard Island white-tailed deer. Tall Timbers Research Station, Tallahassee, Florida, Bulletin 26.
- Rosenberry, C. S., D. R. Diefenbach, and B. D. Wallingford. 2004. Reporting rate variability and precision of white-tailed deer harvest estimates in Pennsylvania. *Journal of Wildlife Management* 68:860-869.
- Severinghaus, C. W. 1949. Tooth development and wear as criteria of age in white-tailed deer. *Journal of Wildlife Management* 13:195-216.
- Skalski, J. R., and J. J. Millspaugh. 2002. Generic variance expressions, precision, and sampling optimization for the sex-age-kill model of population reconstruction. *Journal of Wildlife Management* 66:1308-1316.
- Skalski, J. R., K. E. Ryding, and J. J. Millspaugh. 2005. *Wildlife demography: analysis of sex, age, and count data*. Elsevier Academic Press, Burlington, Massachusetts, USA.
- Stoll, R. J., and W. P. Parker. 1986. Reproductive performance and condition of white-tailed deer in Ohio. *Ohio Journal of Science* 86:164-168.
- Swihart, R. K., H. P. Weeks, Jr., A. L. Easter-Pilcher, and A. J. DeNicola. 1998. Nutritional condition and fertility of white-tailed deer (*Odocoileus virginianus*) from areas with contrasting histories of hunting. *Canadian Journal of Zoology* 76:1932-1941.
- Taylor, T. J. 1996. Condition and reproductive performance of female mule deer in the central Sierra Nevada. *California Fish and Game* 82:122-132.
- Verme, L. J. 1965. Reproductive studies on penned white-tailed deer. *Journal of Wildlife Management* 29:74-79.
- Verme, L. J. 1967. Influence of experimental diets on white-tailed deer reproduction. *Proceedings of the North American Wildlife and Natural Resources Conference* 32:405-420.
- Verme, L. J. 1969. Reproductive patterns of white-tailed deer related to nutritional plane. *Journal of Wildlife Management* 33:881-887.

Table 1. Number of females examined, median conception date for females with conception date data, percent of females bred between 16 October and 16 December, mean embryos per adult female ( $\geq 2$  years of age), and adult female pregnancy rates from 2000 to 2009, Pennsylvania. NOTE: This information is provided for public information, but is not used to make deer management recommendations. WMU level data is used for management recommendations.

<b>Year</b>	<b><i>n</i></b>	<b>Median conception date</b>	<b>Percent bred 16 October to 16 December</b>	<b>Mean embryos per adult female<sup>a</sup></b>	<b>Adult female pregnancy rates (%)<sup>a</sup></b>
2000	1,075	14 November	90	1.60	93
2001	942	17 November	91	1.58	93
2002	520	14 November	86	1.63	93
2003	618	14 November	93	1.59	93
2004	601	15 November	90	1.53	91
2005	883	14 November	90	1.51	92
2006	632	11 November	89	1.54	89
2007 <sup>b</sup>	1,003	16 November	92	1.50	88
2008	1,020	---	---	1.60	93
2009	1,307	---	---	1.60	89

<sup>a</sup> Embryo counts and pregnancy rates adjusted to account for females that were lactating when collected in late spring. As a result of this change, these results may not agree with previous reports.

<sup>b</sup> Final year for conception date research.

Table 2. Number of adult does examined and assessment of deer health by WMU based on samples collected from 2007 to 2009, Pennsylvania.

<b>WMU</b>	<b><i>n</i></b>	<b>Embryos per adult female</b>	<b>Deer health assessment</b>
1A	118	1.58	AT TARGET
1B	68	1.66	AT TARGET
2A	94	1.45	AT TARGET
2B	162	1.67	ABOVE TARGET
2C	141	1.65	ABOVE TARGET
2D	110	1.67	ABOVE TARGET
2E	42	1.67	AT TARGET
2F	117	1.52	AT TARGET
2G	60	1.55	AT TARGET
3A	41	1.68	ABOVE TARGET
3B	81	1.44	AT TARGET
3C	70	1.40	AT TARGET
3D	116	1.29	BELOW TARGET
4A	114	1.51	AT TARGET
4B	76	1.58	AT TARGET
4C	60	1.50	AT TARGET
4D	83	1.57	AT TARGET
4E	69	1.64	AT TARGET
5A	39	1.56	AT TARGET
5B	67	1.70	ABOVE TARGET
5C	93	1.65	AT TARGET
5D	40	1.78	ABOVE TARGET

Table 3. Number of plots sampled, percent with adequate regeneration, and qualitative assessment of forest habitat health by WMU. Data are based on samples collected from 2005 to 2009, Pennsylvania. Results can only be compared to report since 2008 because of changes in methods.

<b>WMU</b>	<b><i>n</i></b>	<b>% plots with adequate regeneration</b>	<b>Forest health assessment</b>
1A	16	48	Fair
1B	20	44	Fair
2A	21	41	Fair
2B	12	47	Fair
2C	45	54	Fair
2D	28	47	Fair
2E	12	39	Fair
2F	37	44	Fair
2G	67	40	POOR
3A	18	58	Fair
3B	47	60	GOOD
3C	26	55	Fair
3D	34	44	Fair
4A	24	57	Fair
4B	25	55	Fair
4C	20	47	Fair
4D	24	37	Fair
4E	14	57	Fair
5A	6	57	Fair
5B	9	33	Fair
5C	12	26	Fair
5D	0	No Data	--

Table 4. Number of deer checked by PGC personnel, number of report cards sent in by successful hunters, and estimated harvests for antlered and antlerless deer by WMU, Pennsylvania, 2009-10.

WMU	Antlered			Antlerless		
	Deer checked	Report cards	Harvest <sup>1</sup>	Deer checked	Report cards	Harvest <sup>a</sup>
1A	317	1,729	5,500	959	3,373	10,700
1B	523	1,660	5,100	1,693	2,695	9,500
2A	357	2,193	6,800	968	3,878	13,900
2B	135	1,579	4,300	481	3,791	20,000
2C	565	2,621	6,500	1,112	3,613	10,900
2D	542	3,270	10,000	1,178	4,995	16,000
2E	251	1,282	3,700	427	1,596	5,300
2F	584	1,819	5,200	898	2,107	6,600
2G	440	2,377	5,200	292	1,579	4,200
3A	334	1,284	3,300	758	2,062	6,000
3B	477	1,760	4,900	1,055	2,657	9,100
3C	544	2,239	6,200	568	2,477	7,100
3D	247	1,360	3,100	560	2,166	6,300
4A	258	1,455	3,700	546	2,168	7,400
4B	309	1,526	4,000	364	1,479	4,100
4C	345	1,892	4,700	748	2,639	7,200
4D	432	1,997	5,000	752	2,559	7,200
4E	388	1,598	4,100	814	2,244	6,300
5A	103	903	2,200	240	1,409	4,200
5B	364	2,715	6,000	1,083	4,250	11,300
5C	405	3,354	7,600	1,238	8,187	23,200
5D	22	589	1,100	163	1,955	3,900
Unk.		49	130		62	190

<sup>a</sup> Estimated harvests are rounded to the nearest 100 or 1,000 based on precision of harvest estimate. Unknown WMU harvests are rounded to the nearest 10 due to the small number.

Table 5. Number of antlered deer aged, age composition of harvests, and approximate number of 2.5-year-old and older males harvested in Pennsylvania, 1997-98 to 2009-10. Three and 4-point antler restrictions started in 2002-03. Percentages may not add up to 100 percent due to rounding.

<b>Year</b>	<b><i>n</i></b>	<b>% 1.5-year-old males</b>	<b>% 2.5-year-old and older males</b>	<b>No. of 2.5-year-old and older males harvested</b>
1997-98	18,563	81	19	33,600
1998-99	21,350	81	19	34,500
1999-00	20,011	80	20	38,900
2000-01	22,145	82	18	36,600
2001-02	18,893	78	22	44,700
2002-03	11,688	68	32	52,900
2003-04	11,367	56	44	62,600
2004-05	10,555	50	50	62,000
2005-06	9,062	52	48	57,800
2006-07	10,819	56	44	59,500
2007-08	8,014	56	44	48,000
2008-09	9,357	52	48	59,200
2009-10	8,443	49	51	55,200

Table 6. Number of antlerless deer aged and age composition of harvests in Pennsylvania, 1997-98 to 2009-10. Percentages may not add up to 100 percent due to rounding.

<b>Year</b>	<b><i>n</i></b>	<b>% 0.5-year-old males</b>	<b>% 0.5-year-old females</b>	<b>% 1.5-year-old and older females</b>
1997-98	28,743	24	20	56
1998-99	24,913	23	20	57
1999-00	18,502	24	20	56
2000-01	30,460	22	20	58
2001-02	25,450	22	18	60
2002-03	30,077	22	18	60
2003-04	28,236	21	18	61
2004-05	24,640	22	18	61
2005-06	19,459	23	19	58
2006-07	19,074	23	19	58
2007-08	17,770	24	20	56
2008-09	17,152	22	18	60
2009-10	16,519	22	18	60

Table 7. Change ( $\lambda^a$ ) in deer density by WMU, 2005 to 2009, Pennsylvania<sup>b</sup>

WMU	2005	2006	2007	2008	2009
1A	1.03	1.00	0.90	0.98	1.04
1B	1.05	1.03	1.01	0.90	0.83
2A	1.01	0.92	0.91	1.14	0.81
2B	1.07	1.01	0.83	0.98	0.94
2C	0.96	0.99	0.93	0.99	0.76
2D	0.96	1.00	0.93	1.16	0.83
2E	1.14	0.90	0.90	1.01	0.87
2F	0.88	0.97	1.00	0.96	0.76
2G	0.95	1.03	0.90	1.14	0.60
3A	0.95	0.92	1.03	0.79	0.99
3B	0.94	0.98	0.96	0.85	1.00
3C	0.92	1.05	1.03	1.06	0.87
3D	0.94	0.99	0.96	1.05	0.62
4A	0.78	1.30	0.98	0.84	0.74
4B	0.83	1.12	0.89	1.13	1.02
4C	0.90	0.94	0.95	1.02	0.86
4D	0.90	1.13	0.88	1.04	0.76
4E	1.08	0.85	1.11	0.94	0.93
5A	0.81	1.05	0.96	1.07	0.98
5B	0.96	0.95	1.03	1.10	0.79
5C	1.03	0.94	1.12	0.91	0.83
5D	0.83	0.93	1.13	1.05	0.88

<sup>a</sup>  $\lambda = 1.00$  indicates no change in deer density. Values greater than 1.00 indicate increases, less than 1.00 indicate decreases.

<sup>b</sup>  $\lambda$ s from prior years may not match  $\lambda$ s previously reported because  $\lambda$ s are updated as new harvest and population data become available.

Table 8. Antlerless license allocations by WMU, 2005-06 to 2010-11, Pennsylvania.

<b>WMU</b>	<b>2005-06</b>	<b>2006-07</b>	<b>2007-08</b>	<b>2008-09</b>	<b>2009-10</b>	<b>2010-11</b>
1A	40,000	42,000	42,000	42,000	42,000	41,705
1B	27,000	30,000	30,000	30,000	30,000	27,844
2A	55,000	55,000	60,000	55,000	55,000	54,879
2B	68,000	68,000	68,000	68,000	68,000	68,000
2C	53,000	49,000	49,000	49,000	49,000	44,107
2D	56,000	56,000	56,000	56,000	56,000	50,123
2E	21,000	21,000	21,000	21,000	21,000	20,407
2F	30,000	28,000	28,000	28,000	28,000	22,148
2G	29,000	19,000	26,000	26,000	26,000	15,210
3A	27,000	29,000	29,000	26,000	26,000	25,247
3B	41,000	43,000	43,000	43,000	43,000	33,761
3C	32,000	27,000	27,000	27,000	27,000	26,358
3D	38,000	38,000	38,000	37,000	37,000	31,622
4A	35,000	29,000	29,000	29,000	29,000	27,521
4B	35,000	31,000	23,000	23,000	23,000	22,148
4C	39,000	39,000	39,000	35,000	35,000	34,351
4D	40,000	40,000	40,000	40,000	40,000	30,052
4E	38,000	38,000	38,000	30,000	30,000	26,899
5A	28,000	25,000	22,000	19,000	19,000	18,269
5B	56,000	53,000	53,000	51,000	51,000	50,812
5C	71,000	79,000	84,000	92,000	113,000	121,960
5D	20,000	20,000	20,000	22,000	22,000	22,000