Annual Deer Population Report & 2020-21 Antlerless License Allocations Recommendations



March 25, 2020

Pennsylvania Game Commission Bureau of Wildlife Management Deer and Elk Section

Summary of 2020-21 Antlerless Allocations to Achieve Deer Plan Goals

2020-21 Deer Plan Recommendation **Deer Plan** 2019-20 **Population Population Approved** 14-day 7-day **WMU Trend** Objective **Allocation** Concurrent Concurrent Comments 49,000 1A Stable Stable 39,000 49,000 Increase harvest by 1.0 deer/mile² to stop 1B Increasing Stable 35,000 35,000 41,000 population growth 2A Stable Stable 46,000 40,000 46,000 43,656 sold in 2019-20 2B Stable Stable 54,000 49,000 45,356 sold in 2019-20 Increase harvest by 1.5 deer/mile² to 2C Stable Reduce 52,000 58,000 63,000 reduce population because of CWD Increase harvest by 1.5 deer/mile² to 2D Stable Reduce 66,000 60,000 77,000 reduce population because of CWD Increase harvest by 1.5 deer/mile² to 2E Stable Reduce 32,000 39,000 39,000 reduce population because of CWD Increase harvest by 1.0 deer/mile² to stop 2F 36,000 **Increasing** Stable 31,000 28,000 population growth 2G Stable Stable 26,000 22,000 27,000 7,000 2H Stable Stable 6,000 6,000 3A Stable Stable 20,000 19,000 21,000 3B 38,000 30,000 33,000 Stable Stable Increase harvest by 1.0 deer/mile² to 46,000 37,000 49,000 3C Stable Reduce reduce deer impact on forests Increase harvest by 1.0 deer/mile² to 3D Stable Reduce 25,000 33,000 36,000 reduce deer impact on forests Increase harvest by 1.5 deer/mile² to 4A Stable Reduce 41,000 49,000 56,000 reduce population because of CWD Increase harvest by 1.5 deer/mile² to Stable Reduce 32,000 33,000 39,000 4B reduce population because of CWD 36,000 28,000 4C Stable Stable 32,000 Increase harvest by 1.0 deer/mile² to 4D Stable Reduce 46,000 45,000 51,000 reduce population because of CWD Increase harvest by 1.0 deer/mile² to 4E Stable Reduce 34,000 36,000 37,000 reduce deer impact on forests Increase harvest by 1.0 deer/mile² to 5A 29,000 Stable Reduce 22,000 26,000 reduce population because of CWD 5B Stable Stable 67,000 58,000 60,000 5C Stable Stable 70,000 70,000 5D Stable Stable 29,000 29,000

Data presented in this report represent collaborative efforts between the U.S. Forest Service, Pennsylvania's Department of Conservation and Natural Resources, the Pennsylvania Cooperative Fish and Wildlife Research Unit at Penn State University, Responsive Management, and the Game Commission's bureaus of Information and Education, Wildlife Habitat Management, and Wildlife Management. For more information on the deer management program and data and methods used to assess progress towards management goals, visit the Game Commission's website, www.pgc.pa.gov, to find the "2009-2018 White-tailed Deer Management Plan".

Deer Management Goals

Deer management goals direct Game Commission staff in formulating deer management recommendations. Current management goals that directly affect antlerless allocations are to manage deer for healthy deer, healthy forest habitat, and acceptable levels of deer-human conflicts. These goals were identified by a group of public stakeholders in 2002 and continue to be supported by a clear majority of Pennsylvania citizens and hunters (Figure 1).




Figure 1. Percent of respondents that agree with deer management goals. The citizen survey was completed by Responsive Management in the fall of 2011, and the deer hunter survey was completed by the Deer and Elk Section and Bureau of Wildlife Management in the fall of 2017.

Step-by-Step Deer Management Recommendation Guide

The deer management program considers data for each goal to arrive at a deer population recommendation in a defined process (see pages 7 and 8). This process has been revised as new data are incorporated into the program and will continue to evolve as more data and understanding are gained. Decision points (i.e., fawn to doe ratio declining?) are based on published protocols from the wildlife and forestry professions.

Do PA residents want fewer or more deer?

This question is answered using results of the survey conducted by Responsive Management of Pennsylvania residents in 2019. If most surveyed residents in a WMU want less deer, the recommendation would be to reduce the deer population. If the deer health goal is met, forest habitat is good, and WMU residents want more deer, the recommendation would be to increase the deer population.

Is CWD present in free-ranging deer?

This question is answered using results from the thousands of deer tested annually for chronic wasting disease (CWD). If CWD is present in free-ranging deer, then management recommendations are to stabilize or reduce WMU populations. Additional antlerless deer can be removed using Deer Management Assistance Program permits in accordance with the CWD response plan. Increasing the antlerless harvest serves 2 purposes that are important to efforts to contain CWD; (1) increased antlerless harvest removes more deer from the population and allows the Game Commission to test more deer in our efforts to obtain the best information on the extent of the disease, and (2) increased antlerless harvest can reduce deer populations and spread of CWD.

Is fawn to doe ratio declining?

This question is answered using results from the age structure of the antlerless harvest. These data are collected each year by trained Game Commission deer agers from across the state. If the proportion of fawns in the antlerless harvest (hereafter referred to as fawn to doe ratio) is declining and the population is not achieving its objective (i.e., population is declining and objective is to maintain a stable deer population), then the antlerless allocation would be reduced to stop the population decline. The antlerless harvest will have the greatest influence on the population because hunting accounts for most deer mortalities in Pennsylvania. If the fawn to doe ratio is stable or if the population is meeting its objective (i.e., population is stable and objective is stable), no management action is taken.

Has deer population been stable or increasing for 6 years?

This question is answered using results from the Pennsylvania Sex-Age-Kill deer population model and deer harvest indices (i.e., antlered harvest, antlerless catch-per-unit-effort). The 6-year time period is necessary because of the 5-year time period to collect the forest data. The sixth

year is added because only 2^{nd} year seedlings are counted in the forest data. As a result, a complete forest data set includes effects of deer from the previous 6 years.

If the deer population is decreasing the recommendation is to stabilize the population at the lower level to see if forest habitat improves given the lower deer population. If the deer population is stable or increasing, the process continues to the next step.

Is forest habitat good?

This question is answered using results from the Pennsylvania Regeneration Study. If 70% of forested plots have adequate regeneration, forest habitat is considered good. If less than 50% of forested plots have adequate regeneration, forest habitat is considered poor. If 50% to 70% of forested plots have adequate regeneration, forest habitat is considered fair.

Is plot to plot regeneration improving?

This question is answered using results from the Pennsylvania Regeneration Study. In this step, results from individual plots are compared in a paired analysis. For example, plot measurements from 2005 are compared to their remeasured results in 2010 to see if regeneration has improved on individual plots. All plots with 2 measures are included in this analysis. If regeneration is improving, then the deer population trend can be stabilized. If regeneration is not improving, the process continues to the next step.

Is plot to plot deer impact improving?

This question is answered using results from the Pennsylvania Regeneration Study. In this step, results from assessments of deer impact on a scale from 1 (very low) to 5 (very high) are compared in the same way as the plot to plot regeneration analysis. If deer impact is improving (i.e., going from a 4 [high] to 3 [moderate]) on enough plots, then the deer population trend can be stabilized. If deer impact is not improving, the process continues to the next step.


Is mean deer impact 3 or less?

This question is answered from the Pennsylvania Regeneration Study. In this step, the mean deer impact for all plots measured in the most recent 5-year period is statistically compared to an objective of 3 (i.e., moderate impact). If deer impact is significantly greater than 3 (moderate), then the deer impact is too high and the deer population should be reduced. If deer impact is less than or not different from 3 (moderate) then the deer population trend can be stabilized.

Guides on pages 7 and 8 are used to develop deer population recommendations based on goals and objectives of deer management plan. Recommendation guide for WMUs 2B, 5C, and 5D differs because of lack of forest data in these highly developed WMUs.




Deer Management Recommendation Process





Deer Management Recommendation Guide

FOR WMUs 2B, 5C, and 5D



Step-by-Step Antlerless License Allocation Calculations


Antlerless allocations are calculated by referring to results from previous seasons. For example, if a population has remained stable with an annual harvest of 3,000, the same level of harvest would be expected to maintain the stable population. If it has taken 3 antlerless licenses to harvest 1 antlerless deer over the last 3 years, the allocation to stabilize this population would be 3,000 antlerless deer harvested x 3 licenses/antlerless deer harvested = 9,000 antlerless licenses.

Table 1. Antlerless licenses needed to harvest 1 antlerless deer (license/deer) based on historic results for each WMU, March 2020.

WMU	2017-18	2018-19	2019-20	3-year Average
1A	4.1	3.8	3.7	3.9
1B	2.7	2.4	2.8	2.6
2A	4.6	4.2	4.4	4.4
2B	3.9	3.8	4.3	4.0
2C	4.0	4.0	4.1	4.0
2D	3.2	3.1	3.8	3.4
2E	3.5	3.1	4.2	3.6
2F	3.3	3.1	3.5	3.3
2G	4.6	4.1	4.3	4.3
2H	3.7	3.3	5.6	4.2
3A	4.0	3.0	3.5	3.5
3B	4.3	3.5	3.7	3.8
3C	3.5	3.1	3.6	3.4
3D	5.9	4.4	5.1	5.1
4A	4.6	5.5	5.8	5.3
4B	3.7	3.9	4.4	4.0
4C	4.5	4.2	4.3	4.3
4D	4.0	3.9	4.5	4.2
4E	3.1	3.4	3.6	3.4
5A	5.7	4.9	4.4	5.0
5B	4.4	4.1	4.5	4.3
5C	4.4	4.2	4.8	4.5
5D	3.9	4.6	4.3	4.3

Trend in Fawn to Doe Ratios, 2014 to 2019

(Supporting data in WMU worksheets, pages 24 to 69)









Forest Regeneration, 2014 to 2018

(Supporting data in WMU worksheets, pages 24 to 69)



Legend

Poor Forest Regeneration Levels Fair Forest Regeneration Levels








Plot to Plot Change in Regeneration, 5-year Change

(Supporting data in WMU worksheets, pages 24 to 69)



Declining Regeneration
No Change in Regeneration


Improving Regeneration





Plot to Plot Change in Deer Impact, 5-year Change

(Supporting data in WMU worksheets, pages 24 to 69)








Deer Impact Level, 2014 to 2018

(Supporting data in WMU worksheets, pages 24 to 69)



Deer Impact is Too High (> 3)


Deer Impact is Acceptable (3 or less)





Post-Hunt Deer Population Trends, 2015 to 2020

(Supporting data in WMU worksheets, pages 24 to 69)



Legend

Declining Deer Population Stable Deer Population Increasing Deer Population








Pennsylvania Residents Opinions on Deer Populations, 2019

(Supporting data in WMU worksheets, pages 24 to 69)



Legend


Most Residents Say Deer Population Too High Most Residents Say Deer Population Just Right Most Residents Say Deer Population Too Low








Residents Opinions on Deer Populations 2011 vs. 2019






Legend


More than 25% say Deer Population Too High Less than 25% say Deer Population Too High and less than 25% say Too Low More than 25% say Deer Population Too Low





Deer Hunters Opinions on Deer Populations 2011 vs. 2017





Legend


More than 25% say Deer Population Too High Less than 25% say Deer Population Too High and less than 25% say Too Low More than 25% say Deer Population Too Low







Chronic Wasting Disease Management Areas, March 2020








2019-20 Regular Firearms Seasons

In 2001, a 12-day concurrent antlered and antlerless firearms season began. The objectives of this longer antlerless season were to give hunters more time to hunt antlerless deer and to create a more consistent harvest from year to year. Beginning in 2008, some WMUs were changed to a 5-day antlered only season followed by a 7-day concurrent antlered and antlerless season. In 2010, 2011, 2014, 2015, and 2017 additional WMUs were changed to the 5/7 season format.



Legend

5-day Antlered Only and 7-day Antlered and Antlerless Concurrent Firearms Season 12-day Antlered and Antlerless Concurrent Firearms Season








2019-20 Antlered Deer Harvest Density

(Antlered deer harvested per square mile of area)



Legend

Less than 2 antlered deer per square mile

2.0 to 3.0 antlered deer per square mile

3.1 to 4.0 antlered deer per square mile

More than 4.0 antlered deer per square mile






Recommendation Guides and Deer Population Datasheets

Recommendation guides (see pages 7 and 8) provide a step-by-step progression through the deer plan goals and measurable objectives to arrive at a deer population recommendation.

Supporting data for these guides are found in the individual WMU datasheets that follow.

WMU Antlerless Allocation Worksheets

Example



2020-21 Pennsylvania Game Commission Antlerless Allocation Worksheet

			VVIV	/io Characteri	Stics	
WMU	1A	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		9%	45%	36%	3%	1.846

2005	Antlered			Deer Harvest Author of University						
2005	Antlered				Antlered Harvest					
		Antlerless	Allocation	Lic/Deer ¹	12,000					
2006	5,468	13,427	40,000	2.9	12,000					
2006	5,791	13,214	42,000	3.1						
2007	4,896	12,490	42,000	3.3	10,000					
2008	5,392	12,611	42,000	3.3						
2009	5,500	10,689	42,000	4.0	8,000					
2010	5,918	11,940	41,705	3.5						
2011	5,171	9,839	42,000	4.3	6,000					
2012	6,078	11,859	42,000	3.5						
2013	6,420	13,915	49,000	3.5						
2014	5,131	10,792	47,000	4.4	4,000					
2015	6,031	9,122	46,000	5.0						
2016	6,500	10,377	46,000	4.4	2,000					
2017	6,279	12,612	52,000	4.1						
2018	5,802	12,442	48,000	3.8	<u> </u>					
2019	6,416	13,160	49,000	3.7	\$\$ \$\$\$ \$\$\$ \$\$\$ \$\$\$ \$\$\$ \$\$\$ \$\$\$ \$\$\$ \$\$\$					

RED=7 day season


POST-HUNT	Deer Population		Trend	Stable
Year	Total			
2008	34,007	160,000		
2009	36,152	140,000		
2010	44,148	110,000		
2011	41,549	120,000		
2012	42,420			
2013	48,472	100,000		
2014	55,114	80,000		
2015	49,169	30,000		
2016	62,237	60,000		
2017	65,707			
2018	53,244	40,000		
2019	46,208	20,000		
2020	51,804	0		
		20° 20° 20° 20° 20° 20° 20° 20° 20° 20°	, Joz. Joz. Joz.	j 2012 2012 2010

DEER HEALTH: Fawn to Doe Ratio²

Trend

Declining

Year	Total
2008	0.44
2009	0.45
2010	0.46
2011	0.43
2012	0.48
2013	0.41
2014	0.46
2015	0.42
2016	0.43
2017	0.40
2018	0.39
2019	0.37




FOREST HEALTH

Plot - Plot Regeneration No change Plot - Plot Deer Impact No change

Regeneration Assessment Fair
ge Mean Deer Impact 3 or less

Year	% Adequate
2003-07	53%
2004-08	54%
2005-09	55%
2006-10	57%
2007-11	55%
2008-12	53%
2009-13	57%
2010-14	55%
2011-15	54%
2012-16	53%
2013-17	50%
2014-18	64%
2012-16 2013-17	53% 50%



Citizen Survey Results 2019 (2011)

Too High 26%(16%) **Just Right** 55%(54%) **Too Low** 13%(23%)

Antlerless Allocation Recommendation							
	Stable	Decrease					
Firearm Season Option	Harvest	Harvest	Harvest				
7 day antlered & 7 day concurrent	57,000	49,000	42,000				
14 day concurrent	46,000	39,000	34,000				

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

2020-21 Pennsylvania Game Commission Antlerless Allocation Worksheet

			VVIV	io characteri	SUCS	
WMU	1B	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		7%	54%	32%	4%	2,115

								Antler	ed Ha	rvest				
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	43.000									
2005	6,382	10,707	27,000	2.5	12,000									
2006	6,773	11,974	30,000	2.5										
2007	6,010	11,400	30,000	2.6	10,000 -									
2008	7,507	13,390	30,000	2.2										
2009	5,089	9,474	30,000	3.2	8,000 -								_	١.
2010	5,470	9,233	27,844	3.0						_	_	_		
2011	6,021	9,508	30,000	3.2	6,000 -									Ш
2012	6,978	11,086	33,000	3.0	0,000			_	. I					
2013	6,835	10,760	31,000	2.9										
2014	5,766	8,788	30,000	3.4	4,000 -									П
2015	6,895	7,671	29,000	3.8										
2016	7,948	8,243	29,000	3.5	2,000 -					-				Н
2017	8,300	13,047	35,000	2.7										
2018	7,971	15,765	37,000	2.4										
2019	8,658	12,738	35,000	2.8		Sq. 50q. 50	51 ₇₀₈	rog to	\$ 9	\$ 3	3 ~	2015 20	6 3	2028

RED=7 day season


ST-HUNT	Deer Population		Trend	Increasing
Year	Total	160,000		
2008	52,810	100,000		
2009	58,926	140,000		
2010	44,469			
2011	46,503	120,000		
2012	51,697	100,000		
2013	55,713	100,000		
2014	53,799	80,000		
2015	47,438			
2016	71,669	60,000		
2017	74,053			
2018	81,376	40,000		
2019	60,756	20,000		
2020	81,659	0		
		20° 20° 20° 20° 20° 20° 20° 20° 20° 20°	or or or or or	or por por s

DEER HEALTH: Fawn to Doe Ratio²

Trend


Stable

Year	Total
2008	0.41
2009	0.43
2010	0.44
2011	0.40
2012	0.37
2013	0.38
2014	0.39
2015	0.40
2016	0.37
2017	0.39
2018	0.35
2019	0.33



FOREST HEALTH Plot Regeneration No change Plot - Plot Deer Impact No change Mean Deer Impact <3

Year	% Adequate
2003-07	41%
2004-08	46%
2005-09	48%
2006-10	54%
2007-11	57%
2008-12	60%
2009-13	55%
2010-14	56%
2011-15	53%
2012-16	48%
2013-17	49%
2014-18	51%



Citizen Survey Results	2019 (2011)	Too High	11%	Just Right	56%	Too Low	26%

Antlerless Allocation Options					
Increase Stable D					
Firearm Season Option	Harvest	Harvest	Harvest		
7 day antlered & 7 day concurrent	41,000	36,000	30,000		
14 day concurrent	35.000	30.000	25.000		

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

2020-21 Pennsylvania Game Commission Antlerless Allocation Worksheet


			VVIV	io characteri	Stics	
WMU	2A	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		7%	61%	29%	3%	1.811

							Antle	red Ha	rvest				
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	12.000								
2005	8,510	19,649	55,000	2.7	12,000 -								
2006	8,104	16,987	55,000	3.2									
2007	6,560	14,322	60,000	3.9	10,000 -								
2008	6,714	15,255	55,000	3.5		_							
2009	6,829	13,920	55,000	4.0	8,000 -								
2010	5,830	13,463	54,879	4.1			_			_			_
2011	7,142	12,677	65,000	4.4	6,000 -								
2012	6,683	12,694	59,000	4.5	0,000					I _			
2013	6,836	13,241	49,000	3.7									
2014	5,131	9,580	46,000	4.8	4,000 -								
2015	6,511	10,507	43,000	4.1									
2016	7,027	9,235	43,000	4.6	2,000 -			11					
2017	6,134	10,866	50,000	4.6									
2018	6,036	10,950	49,000	4.2			▋ੵ▊ੵ			■,■,			
2019	6,929	9,918	46,000	4.4	1	st 20% 2001 20%	\$ 70° 5	20 2012	202 203	20th 25	4 70%	2021 203	2029

RED=7-day season

POST-HUNT Deer Population	Trend	Stable	
---------------------------	-------	--------	--

Year	Total
2008	45,462
2009	50,336
2010	56,286
2011	49,033
2012	68,080
2013	53,996
2014	43,379
2015	30,033
2016	48,723
2017	57,963
2018	46,361
2019	44,587
2020	61,486
2020	61,486




DEER HEALTH: Fawn to Doe Ratio²

Trend

Stable

Year	Total
2008	0.40
2009	0.38
2010	0.36
2011	0.39
2012	0.38
2013	0.39
2014	0.38
2015	0.31
2016	0.36
2017	0.39
2018	0.32
2019	0.32



FOREST HEALTH


Plot - Plot Regeneration No change Plot - Plot Deer Impact No change

Regeneration Assessment Poor

Mean Deer Impact

3 or less

Year % Adequate 2003-07 46% 2004-08 47% 2005-09 46% 2006-10 45% 2007-11 44% 2008-12 42% 2009-13 43% 2010-14 39% 2011-15 43% 2012-16 41% 2013-17 41% 2014-18 31%



Citizen Survey Results 2019 (2011) Too High 25% Just Right 56%	Too Low	13%
--	---------	-----

Antlerless Allocation Options					
	Increase	Stable	Decrease		
Firearm Season Option	Harvest	Harvest	Harvest		
7 day antlered & 7 day concurrent	54,000	46,000	38,000		
14 day concurrent	47,000	40,000	33,000		

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

2020-21 Pennsylvania Game Commission Antlerless Allocation Worksheet

		WIVIO Characteristic				
WMU	2B	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		30%	44%	21%	0%	1,363

Deer Harvest Antlered Harvest Antlered Antlerless Allocation Lic/Deer¹ Year 12,000 2005 4.4 5,182 14,459 68,000 2006 5,759 16,505 68,000 3.9 10,000 2007 4,372 15,332 68,000 3.9 2008 3,964 15,251 68,000 4.1 2009 4,297 19,866 68,000 3.3 8,000 3,976 13,008 68,000 2010 4.8 2011 4,472 16,550 71,000 3.6 6,000 2012 4,837 15,955 67,000 3.8 2013 5,610 14,389 62,000 4.3 4,000 2014 4,267 13,165 60,000 4.5 2015 5,191 15,379 61,000 3.9 2,000 2016 5,801 14,317 60,000 4.2 2017 4,458 13,930 60,000 3.9 2018 5,036 12,318 58,000 3.8 2019 5,503 10,374 54,000 4.3

POST-HUNT Deer Population	Trend	Stable


Year	Total	
2008		
2009		
2010	Harvest indi	ces (i.e., antlered harvest, antlerless lic/deer), not PASAK model, used to monitor population trend
2011		
2012		
2013		
2014		

DEER HEALTH: Fawn to Doe Ratio²

Trend

Stable

Year	Total
2008	0.43
2009	0.48
2010	0.47
2011	0.41
2012	0.45
2013	0.44
2014	0.44
2015	0.37
2016	0.41
2017	0.45
2018	0.36
2019	0.37



FOREST HEALTH Plot - Plot Regeneration Plot - Plot Deer Impact Regeneration Assessment Mean Deer Impact

Year	% Adequate	
2003-07		
2004-08		
2005-09		Forest data not considered in this developed WI
2006-10		
2007-11		
2008-12		
2009-13		

Citizen Survey Results	2019 (2011)	Too High	38%(32%) Just Right	52%	Too Low	9%

Antlerless Allocation Options						
	Increase	Stable	Decrease			
Firearm Season Option	Harvest	Harvest	Harvest			
14 day concurrent	54,000	49,000	44,000			

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

2020-21 Pennsylvania Game Commission Antlerless Allocation Worksheet


			VVIV	no Characteri	Stics	
WMU	2C	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		6%	68%	24%	10%	2.934

er Harve	est														
							Α	ntlere	d Ha	rvest					
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	13.000										
2005	7,413	13,683	53,000	3.8	12,000 -										
2006	9,049	12,094	49,000	4.0											
2007	8,441	11,619	49,000	4.1	10,000 -										
2008	7,476	12,752	49,000	3.8					_						
2009	6,508	10,870	49,000	4.5	8,000 -									₽	Н
2010	8,528	9,579	44,107	4.6											
2011	8,249	12,793	58,000	4.5	6,000 -		ш				Ш	ш		Ш	Ш
2012	7,600	10,822	50,000	4.6	,,,,,,										
2013	7,219	10,957	43,000	3.9	4.000										
2014	7,016	8,985	38,000	4.5	4,000 -										П
2015	9,134	7,269	31,000	4.3											
2016	8,300	6,869	31,000	4.6	2,000 -										Н
2017	9,792	7,724	31,000	4.0											
2018	9,572	11,134	44,000	4.0			┸,		_			⊢ ■			
2019	9,426	12,743	52,000	4.1	า์	it in initial	100° 1	90° 90'E	, 2017	202 20	3 20%	2015	20° 20°	3 203	2029

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	87,046
2009	72,402
2010	62,340
2011	66,729
2012	64,888
2013	61,386
2014	68,683
2015	66,027
2016	83,350
2017	69,034
2018	113,659
2019	86,087
2020	97,246




DEER HEALTH: Fawn to Doe Ratio²

Approximately 50% of WMU is a CWD DMA in 2019

Trend

Stable

Year	Total
2008	0.43
2009	0.43
2010	0.40
2011	0.40
2012	0.38
2013	0.44
2014	0.42
2015	0.37
2016	0.35
2017	0.37
2018	0.34
2019	0.35



FOREST HEALTH


Plot - Plot Regeneration No change Plot - Plot Deer Impact Increasing

Regeneration Assessment

Fair

Mean Deer Impact 3 or less

Year	% Adequate
2003-07	54%
2004-08	59%
2005-09	58%
2006-10	58%
2007-11	59%
2008-12	56%
2009-13	57%
2010-14	58%
2011-15	62%
2012-16	63%
2013-17	60%
2014-18	58%



Citizen Survey Results 2019 (2011)

Too High

19%(13%) Just Right

52%(50%) **Too Low**

23%(26%)

Antlerless Allocation Options							
Firearm Season Option	Increase Harvest	Stable Harvest	Decrease Harvest				
7 day antlered & 7 day concurrent	63,000	45,000	34,000				
14 day concurrent	58,000	41,000	31,000				

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.

2020-21 Pennsylvania Game Commission Antlerless Allocation Worksheet


			VVIV	/io Characteri	Stics	
WMU	2D	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		5%	60%	31%	2%	2.486

Deer Harvest							
					Antlered Harvest		
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	14.000		
2005	9,975	22,054	56,000	2.5	14,000		
2006	10,896	20,437	56,000	2.7			
2007	9,118	18,099	56,000	3.1	12,000		
2008	9,508	15,591	56,000	3.5			
2009	9,977	15,962	56,000	3.5	10,000		
2010	11,540	18,046	50,123	2.8			
2011	11,130	19,257	60,000	3.1	8,000		
2012	13,660	20,839	62,000	3.0			
2013	13,704	21,614	61,000	2.8			
2014	11,417	16,441	61,000	3.7	6,000		
2015	12,292	15,728	55,000	3.5			
2016	12,843	16,447	55,000	3.3	4,000		
2017	14,716	17,033	55,000	3.2			
2018	11,847	20,345	63,000	3.1	2,000		
2019	12,971	17,472	66,000	3.8	\$ \$		

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	69,732
2009	88,666
2010	86,493
2011	101,182
2012	102,440
2013	113,774
2014	144,084
2015	110,214
2016	117,823
2017	112,499
2018	140,281
2019	105,280
2020	114,679




DEER HEALTH: Fawn to Doe Ratio²

Approximately 13% of WMU is a CWD DMA in 2019

Trend

Declining

Year	Total
2008	0.44
2009	0.39
2010	0.42
2011	0.44
2012	0.42
2013	0.42
2014	0.40
2015	0.37
2016	0.36
2017	0.37
2018	0.34
2019	0.33




FOREST HEALTH

Plot - Plot Regeneration No change Plot - Plot Deer Impact No change

Fair

3 or less

Year	% Adequate
2003-07	52%
2004-08	54%
2005-09	51%
2006-10	52%
2007-11	49%
2008-12	46%
2009-13	50%
2010-14	45%
2011-15	44%
2012-16	50%
2013-17	48%
2014-18	41%



Citizen Survey Results 2019 (2011)

Too High

26%(23%) Just Right

57%(52%) **Too Low**

Regeneration Assessment

13%(19%)

Antlerless Allocation Options							
Firearm Season Option	Increase Harvest	Stable Harvest	Decrease Harvest				
7 day antlered & 7 day concurrent	77,000	64,000	56,000				
14 day concurrent	60,000	50,000	44,000				

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

2020-21 Pennsylvania Game Commission Antlerless Allocation Worksheet


			WIVIO Characteristics					
WMU	2E	% Developed	d % Forest	%Ag/Field	%Public	Area (sq mi)		
		5%	65%	26%	6%	1.427		

								Antle	red H	arvest					
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	12.000										
2005	4,093	7,471	21,000	2.8	12,000										
2006	5,358	7,360	21,000	2.8											
2007	3,642	6,398	21,000	3.2	10,000										
2008	4,984	6,179	21,000	3.3											
2009	3,673	5,298	21,000	4.0	8,000										
2010	4,178	5,952	20,407	3.5										_	
2011	4,116	7,073	25,000	3.5	6,000									Ш	
2012	4,785	5,561	21,000	3.8			_	_					_		ı
2013	4,883	7,973	22,000	2.8	4.000					₋▮		_			
2014	4,440	5,593	21,000	3.8	4,000										
2015	4,742	5,263	21,000	4.0											
2016	5,221	5,215	21,000	4.1	2,000					Н					Н
2017	6,929	6,214	22,000	3.5											
2018	6,274	8,693	27,000	3.1	_ +								, ■,		_
2019	6,370	7,641	32,000	4.2	\$	\$ 706 V	51 ₂₀₀ 0	SO (\$ \$	102 Y	03° 00	\$ \$	2026 25	37 29	ه ک

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	32,623
2009	42,709
2010	38,317
2011	38,134
2012	30,384
2013	44,546
2014	45,529
2015	50,549
2016	43,081
2017	43,144
2018	56,635
2019	47,171
2020	62,753




Approximately 51% of WMU is a CWD DMA in 2019

Trend

Stable

Year	Total
2008	0.40
2009	0.37
2010	0.46
2011	0.41
2012	0.43
2013	0.40
2014	0.36
2015	0.36
2016	0.36
2017	0.33
2018	0.31
2019	0.33




FOREST HEALTH

Plot - Plot Regeneration No change Plot - Plot Deer Impact No change

Regeneration Assessment Fai

Fair 3 or less

Year	% Adequate
2003-07	53%
2004-08	50%
2005-09	47%
2006-10	50%
2007-11	52%
2008-12	52%
2009-13	56%
2010-14	61%
2011-15	63%
2012-16	56%
2013-17	60%
2014-18	56%



Citizen Survey Results 2019 (2011)

Too High

20%(13%) Just Right

56%(48%) **Too Low**

22%(31%)

Antierless Allo	cation Option	ons	
Firearm Season Option	Increase	Stable	Decrease
	Harvest	Harvest	Harvest
7 day antlered & 7 day concurrent	39,000	31,000	26,000
14 day concurrent	39,000	31,000	26,000

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VV IV	io characteri	Stics	
WMU	2F	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		2%	88%	7%	56%	2,409

	est			-	Antlere	d Han	/est							
Year Antlered Antlerless Allocation Lic/Deer ¹			12.000											
2005	6,013	8,322	30,000	3.5	12,000									
2006	7,153	8,030	28,000	3.5										
2007	4,795	7,132	28,000	3.9	10,000									
2008	6,990	9,117	28,000	3.0										
2009	5,167	6,648	28,000	4.3	8,000									_
2010	6,403	5,657	22,148	4.0			_					_		
2011	5,393	6,737	34,000	5.0	6,000									
2012	7,139	6,067	27,000	4.5	0,000			_						
2013	6,607	8,008	29,000	3.6	4.000									
2014	5,979	5,915	27,000	4.6	4,000									
2015	6,989	5,434	22,000	4.1										
2016	7,678	6,718	22,000	3.3	2,000									1
2017	9,489	7,200	24,000	3.3										
2018	7,665	7,533	23,000	3.1	_ +			_ 📕 _	■,■,		₽			
2019	9,014	8,816	31,000	3.5	26	5 706 70°	1 20gs	ing Jose	2017	25 20°S	, 20 ⁷ 2	25 JO 26	2027	29° 20'

RED=7-day season

POST-HUNT Deer Population	Trend	Increasing

Year	Total
2008	47,288
2009	67,724
2010	46,887
2011	70,765
2012	53,210
2013	83,063
2014	65,614
2015	61,020
2016	67,152
2017	74,387
2018	108,575
2019	87,309
2020	98,104
2016 2017 2018 2019	67,152 74,387 108,575 87,309




Approximately 8% of WMU is a CWD DMA in 2019

Trend

Stable

Year	Total
2008	0.35
2009	0.34
2010	0.39
2011	0.40
2012	0.39
2013	0.38
2014	0.37
2015	0.34
2016	0.33
2017	0.36
2018	0.37
2019	0.28




FOREST HEALTH

Plot - Plot Regeneration No change Plot - Plot Deer Impact No change

Regeneration Assessment Good

3 or less

Year	% Adequate
2003-07	47%
2004-08	50%
2005-09	50%
2006-10	54%
2007-11	54%
2008-12	54%
2009-13	54%
2010-14	58%
2011-15	61%
2012-16	65%
2013-17	69%
2014-18	71%



Citizen Survey Results 2019 (2011)

Too High 19%(10%) **Just Right** 48%(39%) **Too Low** 26%(42%)

Antierless Allo	cation Option	ons	
Firearm Season Option	Increase	Stable	Decrease
	Harvest	Harvest	Harvest
7 day antlered & 7 day concurrent	36,000	28,000	20,000
14 day concurrent	28,000	22,000	16,000

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	10 Characteri	Stits	
WMU	2G	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		4%	82%	7%	57%	3,117

						Ar	tlered Ha	rvest				
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	12.000							
2005					12,000							
2006												
2007					10,000 -							_
2008												
2009	3,802	1,046			8,000 -							-
2010	5,088	2,627										
2011	4,957	4,117			6,000 -							
2012	4,976	4,915			5,555		_					
2013	5,018	6,881	28,000	4.1	4.000							
2014	4,839	4,671	22,000	4.7	4,000							
2015	6,073	4,143	22,000	5.4								
2016	6,201	3,996	21,000	5.3	2,000 -				\blacksquare			_
2017	8,193	5,516	25,500	4.6								
2018	6,296	7,372	30,000	4.1								_
2019	8,062	6,123	26,000	4.3		in hay hay hay ha	9 29 27	\$ 3	N N	Nº X	1 29	%

RED=7-day season


Deer Population Trend Stable	Deer Population Tren	d Stable
-------------------------------------	----------------------	----------

Year	Total
2008	
2009	55,234
2010	41,008
2011	45,743
2012	55,997
2013	57,014
2014	49,313
2015	40,343
2016	65,521
2017	67,942
2018	81,757
2019	55,221
2020	70,946



Trend Stable

Year	Total
2008	0.34
2009	0.35
2010	0.39
2011	0.35
2012	0.25
2013	0.35
2014	0.30
2015	0.34
2016	0.31
2017	0.29
2018	0.35
2019	0.26



FOREST HEALTH


Plot - Plot Regeneration No change Plot - Plot Deer Impact Increasing

Regeneration Assessment

Fair

Mean Deer Impact 3 or less

% Adequate
53%
53%
54%
54%
55%
56%
55%
54%
56%
52%
55%
52%



Citizen Survey Results 2019 (2011)

Too High

13%(3%) **Just Right**

49%(39%) **Too Low**

35%(55%)

Antlerless Allo	cation Option	ons	
	Increase	Stable	Decrease
Firearm Season Option	Harvest	Stable Decrease Harvest Harvest 27,000 14,000	
7 day antlered & 7 day concurrent	41,000	27,000	14,000
14 day concurrent	34,000	22,000	12,000

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.

			VVIV	10 Characteri	Stits	
WMU	2H	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		4%	86%	6%	27%	1,001

						Antlered Harvest
Year	Antlered	Antierless	Allocation	Lic/Deer ¹		, and creating rest
2005	7	7	7	.,	12,000	
2006						
2007					10,000	+
2008						
2009	1,471	1,046			8,000	
2010	1,670	990				
2011	1,323	1,321			6,000	
2012	1,565	1,459			0,000	
2013	1,475	1,657	6,000	3.7	4.000	
2014	1,670	1,064	5,500	5.2	4,000	
2015	1,426	1,419	6,500	4.6		
2016	1,867	1,861	6,000	3.2	2,000	
2017	1,726	1,889	7,000	3.7		
2018	2,478	1,812	6,000	3.3	-	
2019	2,404	1,086	6,000	5.6		\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\


RED=7-day season

Deer Popul	Year Total 2008 19,730 2010 11,565 2011 18,952 2012 13,917 2013 16,895 2014 16,537 2015 16,872 2016 15,430 2017 15,704 2018 38,649 2019 18,919		Trend	Stable
Year	Total	450,000		
2008		160,000		
2009	19,730	140,000		
2010	11,565	7,11		
2011	18,952	120,000		
2012	13,917			
2013	16,895	100,000		
2014	16,537	80,000		
2015	16,872	30,000		
2016	15,430	60,000		
2017	15,704			
2018	38,649	40,000		
2019	18,919	20,000		
2020	25,314	20,000		
		0	<u> </u>	
		208 208 202 202 2	or sor sor sor sor s	n 10 10 10 10 10 10 10 10 10 10 10 10 10

Trend

Stable

Year	Total
2008	0.43
2009	0.40
2010	0.35
2011	0.33
2012	0.39
2013	0.23
2014	0.33
2015	0.33
2016	0.24
2017	0.40
2018	0.35
2019	0.32



FOREST HEALTH


Plot - Plot Regeneration No change Plot - Plot Deer Impact No change*

Regeneration Assessment

Fair <3

Mean Deer Impact

Year % Adequate 2003-07 44% 2004-08 43% 2005-09 47% 2006-10 52% 2007-11 51% 2008-12 51% 2009-13 57% 2010-14 56% 2011-15 57% 55% 2012-16 2013-17 62% 2014-18 54%



Citizen Survey Results 2019 (2011)

Too High

13%(3%) Just Right

49%(39%) **Too Low**

35%(55%)

Antlerless Allo	cation Option	ons	
Firearm Season Option	Increase	Stable	Decrease
	Harvest	Harvest	Harvest
7 day antlered & 7 day concurrent	11,000	7,000	3,000
14 day concurrent	9,000	6,000	2,000

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	/io Characteri	Stics	
WMU	3A	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		2%	78%	17%	10%	1.506

							Antle	ered Ha	rvest					
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	12.000									
2005	3,981	8,657	27,000	3.1	12,000									
2006	4,527	8,818	29,000	3.2										
2007	3,359	7,803	29,000	3.6	10,000									_
2008	4,132	7,478	26,000	3.4										
2009	3,310	5,998	26,000	4.4	8,000									
2010	3,751	6,469	25,247	3.9										
2011	3,345	6,672	26,000	3.9	6,000									
2012	4,278	6,673	26,000	3.9	5,555									
2013	4,177	5,430	23,000	4.2	4 000		_		_	_	_			1
2014	3,308	4,253	18,000	4.2	4,000			I _		I _				
2015	4,314	4,005	19,000	4.8										
2016	5,432	3,776	15,000	4.0	2,000									Н
2017	5,419	5,014	20,000	4.0										
2018	4,825	7,430	22,000	3.0	_ +				-		↓ ■			<u>_</u>
2019	5,704	5,663	20,000	3.5		or gor gol go	30,700,	00,00	00° 00	3 04	2015 1	26° 50	1 20%	2

RED=7-day season

POST-HUNT Deer Population	Trend	Stable


Year	Total
2008	32,425
2009	32,513
2010	31,412
2011	39,532
2012	31,224
2013	41,358
2014	45,317
2015	36,181
2016	49,307
2017	49,426
2018	55,441
2019	39,832
2020	54,040



Trend

Stable

Year	Total
2008	0.33
2009	0.35
2010	0.42
2011	0.36
2012	0.34
2013	0.36
2014	0.36
2015	0.36
2016	0.27
2017	0.32
2018	0.33
2019	0.32




FOREST HEALTH

Plot - Plot Regeneration No change Plot - Plot Deer Impact No change

Regeneration Assessment F

Fair 3 or less

Year % Adequate 2003-07 65% 2004-08 63% 2005-09 62% 2006-10 61% 2007-11 63% 2008-12 60% 2009-13 66% 2010-14 66% 2011-15 67% 65% 2012-16 2013-17 69% 2014-18 64%



Citizen Survey Results 2019

Too High

18(3%) Ju

Just Right

57%(32%) **Too Low**

21%(59%)

Antlerless Allocation Options					
Increase Stable Decrease Firearm Season Option Harvest Harvest Harvest					
7 day antlered & 7 day concurrent 14 day concurrent	26,000 24,000	21,000 19,000	16,000 15,000		

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	io characteri	Stits	
WMU	3B	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		6%	79%	11%	21%	2,218

					Antlered Harvest	
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	42.000	
2005	5,980	10,871	41,000	3.7	12,000	
2006	6,530	10,563	43,000	4.0		
2007	5,933	10,177	43,000	4.2	10,000	
2008	5,469	9,857	43,000	4.3		
2009	4,865	9,112	43,000	4.7	8,000	
2010	5,369	7,585	33,761	4.5		_ [[] _ [
2011	5,935	7,707	40,000	5.2	6,000	
2012	5,752	8,701	40,000	4.6		
2013	6,153	8,718	39,000	4.5	4.000	
2014	6,039	8,055	33,000	4.1	4,000	
2015	6,840	7,359	28,000	3.8		
2016	7,481	7,290	28,000	3.8	2,000	
2017	8,945	6,970	30,000	4.3		
2018	6,977	8,354	29,000	3.5	-	
2019	7,558	10,264	38,000	3.7	Son	2012 2012 2012 2013 2013 2013

RED=7-day season


Deer Population	Trend	Stable
-----------------	-------	--------

Year	Total
2008	56,162
2009	46,869
2010	48,895
2011	49,768
2012	58,481
2013	53,709
2014	63,803
2015	55,249
2016	76,808
2017	80,598
2018	76,249
2019	51,976
2020	62,489



Trend Declining

Year	Total
2008	0.37
2009	0.40
2010	0.40
2011	0.38
2012	0.39
2013	0.39
2014	0.38
2015	0.38
2016	0.37
2017	0.33
2018	0.31
2019	0.30




FOREST HEALTH

Plot - Plot Regeneration No change Plot - Plot Deer Impact No change

Regeneration Assessment Good ge Mean Deer Impact 3 or less

Year % Adequate 2003-07 58% 2004-08 59% 2005-09 62% 2006-10 62% 2007-11 60% 2008-12 65% 2009-13 67% 2010-14 65% 2011-15 61% 2012-16 64% 2013-17 57% 2014-18 63%



Citizen Surve	y Results	2019	(2011)	1

Too High 20%(7%) **Just Right** 55%(59%) **Too Low** 17%(24%)

Antlerless Allocation Options					
Increase Stable Decrease					
Firearm Season Option	Harvest	Harvest	Harvest		
7 day antlered & 7 day concurrent	41,000	33,000	24,000		
14 day concurrent	38,000	30,000	22,000		

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	/io Characteri	Stics	
WMU	3C	% Developed	d % Forest	%Ag/Field	%Public	Area (sq mi)
		4%	75%	16%	3%	2.187

								Antlei	ed Ha	rvest				
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	12.000									
2005	5,821	11,198	32,000	2.8	12,000 -									
2006	6,673	9,248	27,000	2.9										
2007	5,278	9,586	27,000	2.8	10,000 -									
2008	6,288	7,258	27,000	3.7								i		
2009	6,196	7,084	27,000	3.9	8,000 -					_				
2010	6,211	8,309	26,358	3.2							_			
2011	7,103	9,943	29,000	2.9	6,000 -		_							
2012	7,854	10,508	35,000	3.3	0,000		. I							
2013	7,004	12,683	35,000	2.8	4.000									
2014	6,526	10,302	32,000	3.1	4,000 -									
2015	7,614	10,460	36,000	3.4										
2016	8,629	10,968	36,000	3.3	2,000 -					-				
2017	8,703	11,860	42,000	3.5										
2018	7,739	12,172	38,000	3.1			▋╷▊				▋Ţ▐			
2019	9,382	12,808	46,000	3.6	1	igo fago fag	2) ²⁰ 08	2009 25'	202	202 20	3 2014	105 70°	6 20 ² 1	203º 203

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	45,511
2009	54,141
2010	65,624
2011	59,245
2012	64,359
2013	67,720
2014	58,925
2015	67,997
2016	83,206
2017	85,083
2018	79,925
2019	57,169
2020	75,360




Fair

>3

DEER HEALTH: Fawn to Doe Ratio²

Trend Declining


Year	Total
2008	0.31
2009	0.35
2010	0.32
2011	0.34
2012	0.32
2013	0.35
2014	0.36
2015	0.34
2016	0.38
2017	0.33
2018	0.32
2019	0.27




FOREST HEALTH

 Plot - Plot Regeneration
 No change
 Plot - Plot Deer Impact
 No change
 Mean Deer Impact

 . lot negeneration	ito change		oc beer impace	TTO CITATI
	1000	,		
	1009	6 ⊤		





Citizen Survey Results 2019 (2011)

Too High 30%(10%) **Just Right** 55%(61%) **Too Low** 11%(20%)

Antlerless Allocation Options					
Firearm Season Option	Increase Harvest	Stable Harvest	Decrease Harvest		
7 day antlered & 7 day concurrent	49,000	42,000	34,000		
14 day concurrent	37,000	32,000	26,000		

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	io Characteri	Stics	
WMU	3D	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		11%	74%	6%	16%	2.101

							An	tlered	d Harv	est					
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	13.000										
2005	3,865	7,254	38,000	5.1	12,000										
2006	4,969	7,445	38,000	5.0											
2007	3,647	7,017	38,000	5.3	10,000										
2008	3,899	6,925	37,000	5.3											
2009	3,096	6,265	37,000	5.9	8,000										
2010	3,884	5,509	31,622	5.8											
2011	4,509	7,163	39,000	5.4	6,000										
2012	4,039	6,010	39,000	6.5											
2013	3,446	4,986	32,000	6.4	4 000								_		
2014	4,155	5,203	25,000	4.8	4,000					Ι.					
2015	3,500	3,655	25,000	6.9											
2016	4,272	4,235	25,000	5.9	2,000										
2017	4,656	4,187	25,000	5.9											
2018	5,189	5,690	25,000	4.4	_ +			┻┯┻	, I		_	⊢ ■	,		_
2019	6,016	4,932	25,000	5.1	\$	\$ 206 2001	200° 201	8,00	0 ¹	303	2014	2015 Y	50% V	5,05	

RED=7-day season


	POST-HUNT Deer Population	Trend	Stable
--	---------------------------	-------	--------

Year	Total
2008	31,623
2009	37,563
2010	25,378
2011	30,250
2012	31,299
2013	29,225
2014	25,127
2015	33,778
2016	28,957
2017	33,302
2018	30,727
2019	33,798
2020	48,663
	•



Trend Stable

Year	Total
2008	0.32
2009	0.35
2010	0.31
2011	0.34
2012	0.34
2013	0.30
2014	0.29
2015	0.27
2016	0.31
2017	0.24
2018	0.30
2019	0.27




FOREST HEALTH

Plot - Plot Regeneration No change Plot - Plot Deer Impact No change

Regeneration Assessment Mean Deer Impact

Fair >3

Year	% Adequate
2003-07	56%
2004-08	54%
2005-09	55%
2006-10	58%
2007-11	57%
2008-12	59%
2009-13	61%
2010-14	61%
2011-15	57%
2012-16	63%
2013-17	57%
2014-18	59%



Citizen Survey Results 2019 (2011)

Too High 30%(13%) Just Right

52%(57%) **Too Low**

13%(24%)

Antlerless Allocation Options							
Increase Stable Decrease Firearm Season Option Harvest Harvest Harvest							
7 day antlered & 7 day concurrent	36,000	25,000	15,000				
14 day concurrent	33,000	23,000	14,000				

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.


 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

			WIVIO Characteristics					
WMU	4A	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)		
		4%	70%	24%	15%	1.736		

er Harve	st					
						Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	12.000	
2005	3,714	7,578	35,000	4.5	12,000	
2006	5,871	7,827	29,000	3.6		
2007	4,477	6,735	29,000	4.2	10,000 -	
2008	4,187	6,874	29,000	4.2		
2009	3,733	7,414	29,000	3.9	8,000 -	
2010	3,761	6,401	27,521	4.3		
2011	4,849	6,527	28,000	4.3	6,000 -	
2012	4,245	6,463	29,000	4.5	0,000	
2013	4,961	5,981	28,000	4.7		
2014	3,317	6,802	28,000	5.6	4,000 -	
2015	5,095	6,360	30,000	4.7		
2016	4,423	5,726	30,000	5.2	2,000 -	
2017	4,810	6,475	30,000	4.6		
2018	5,142	6,395	38,000	5.5		│█▗▊▗▊▗▊▗▊▗▊▗▊▗▊▗▊▗▊▗▊▗▊▗▊ ▗▊
2019	5,981	5,250	41,000	5.8		à tà tà tà tà tà tà tà, tà, tà, tà, tà,


RED=7-day season

Year	Total
2008	47,414
2009	34,628
2010	30,789
2011	38,125
2012	49,191
2013	36,579
2014	42,196
2015	23,772
2016	48,538
2017	29,746
2018	39,238
2019	42,174
2020	47,047




DEER HEALTH: Fawn to Doe Ratio ² 100% of WMU is a CWD DMA in 2019	Trend S	table
--	---------	-------

Year	Total
2008	0.34
2009	0.34
2010	0.32
2011	0.38
2012	0.32
2013	0.30
2014	0.32
2015	0.37
2016	0.29
2017	0.30
2018	0.30
2019	0.30



FOREST HEALTH		Reg	generation Assessment	Good	
	Plot - Plot Regeneration Declining	Plot - Plot Deer Impact No change	Mean Deer Impact	<3	

Year	% Adequate
2003-07	60%
2004-08	64%
2005-09	64%
2006-10	61%
2007-11	63%
2008-12	60%
2009-13	59%
2010-14	61%
2011-15	63%
2012-16	67%
2013-17	68%
2014-18	75%



Citizen Survey Results	2019 (2011)	Too High	14%(4%)	Just Right	45%(45%)	Too Low	37%(42%)

Antlerless Allocation Options							
	Increase	Stable	Decrease				
Firearm Season Option	Harvest	Harvest	Harvest				
7 day antlered & 7 day concurrent	56,000	42,000	33,000				
14 day concurrent	49,000	37,000	29,000				

 $^{^{1}}$ - The number of antierless licenses sold that it takes to harvest an antierless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	io Characteri	Stics	
WMU	4B	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		6%	65%	27%	15%	1.591

						Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	12.000	
2005	3,571	6,644	35,000	5.2	12,000	
2006	5,026	6,626	31,000	4.6		
2007	3,472	4,509	23,000	5.0	10,000	
2008	3,917	3,846	23,000	5.9		
2009	4,011	4,061	23,000	5.7	8,000	
2010	4,458	5,113	22,148	4.4		
2011	5,341	5,498	23,000	4.2	6,000	
2012	5,622	5,636	26,000	4.6	5,555	
2013	5,312	5,769	24,000	4.2	4.000	
2014	4,611	5,630	26,000	4.6	4,000	
2015	5,701	6,961	26,000	3.8		
2016	5,164	6,151	26,000	4.2	2,000	
2017	5,602	7,061	26,000	3.7		
2018	5,273	6,757	26,000	3.9		
2019	5,722	7,305	32,000	4.4		¢ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$

RED=7-day season


POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	30,479
2009	39,044
2010	43,550
2011	37,273
2012	60,340
2013	52,903
2014	50,517
2015	45,362
2016	57,846
2017	55,941
2018	52,407
2019	50,252
2020	54,044




DEER HEALTH: Fawn to Doe Ratio ²	100% of WMU is a CWD DMA in 2019	Trend	Declining
			200

Year	Total
2008	0.38
2009	0.44
2010	0.38
2011	0.41
2012	0.33
2013	0.31
2014	0.40
2015	0.36
2016	0.35
2017	0.34
2018	0.30
2019	0.29



FOREST HEALTH			Rege	neration Assessment	Fair
POREST HEALTH	Plot - Plot Regeneration No change	Plot - Plot Deer Impact	No change	Mean Deer Impact	3 or less

Year	% Adequate
2003-07	61%
2004-08	60%
2005-09	58%
2006-10	60%
2007-11	64%
2008-12	61%
2009-13	59%
2010-14	60%
2011-15	63%
2012-16	68%
2013-17	59%
2014-18	57%



Citizen Survey Results 2019 (2011) Too High 16%(6%) Just Right 53%(53%) Too Low 21%	Citizen Survey Results	019 (2011)	Too High 16%(6%	6) Just Right	53%(53%) 100 LOW	21%(33%)
---	------------------------	------------	------------------------	---------------	-------------------------	----------

Antlerless Allo	cation Option	ons	
Firearm Season Option	Increase Harvest	Stable Harvest	Decrease Harvest
7 day antlered & 7 day concurrent	39,000	29,000	23,000
14 day concurrent	33,000	25,000	20,000

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $^{^{2}}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antierless harvest.


			VVIV	io characteri	Stics	
WMU	4C	% Developed	d % Forest	%Ag/Field	%Public	Area (sq mi)
		8%	71%	17%	15%	1.717

er Harve	est					
						Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	12.000	
2005	5,891	9,805	39,000	3.9	12,000 -	
2006	6,115	8,883	39,000	4.2		
2007	4,828	9,375	39,000	4.1	10,000 -	
2008	5,015	8,027	35,000	4.3		
2009	4,745	7,163	35,000	4.9	8,000 -	
2010	5,724	8,357	34,351	4.2		_
2011	5,525	7,392	35,000	4.7	6,000 -	
2012	5,335	7,823	35,000	4.5	0,000	
2013	5,180	6,922	27,000	3.9		
2014	4,830	4,996	25,000	5.1	4,000 -	
2015	5,381	4,976	25,000	5.1		
2016	6,381	5,273	25,000	4.8	2,000 -	
2017	6,799	6,464	29,000	4.5		
2018	5,781	7,155	30,000	4.2		
2019	6,975	8,328	36,000	4.3		e to

RED=7-day season

POST-HUNT Deer Population	Trend	Stable


Year	Total
2008	44,569
2009	45,224
2010	44,256
2011	58,091
2012	45,093
2013	45,586
2014	49,072
2015	50,265
2016	55,068
2017	55,311
2018	61,317
2019	55,122
2020	55,238



Trend

Stable

Year	Total
2008	0.42
2009	0.41
2010	0.42
2011	0.38
2012	0.39
2013	0.40
2014	0.40
2015	0.37
2016	0.36
2017	0.37
2018	0.37
2019	0.33



FOREST HEALTH


Regeneration Assessment

Fair

Plot - Plot Regeneration No change Plot - Plot Deer Impact No change

Mean Deer Impact 3 or less

Year % Adequate 2003-07 66% 2004-08 63% 2005-09 63% 2006-10 63% 2007-11 60% 2008-12 61% 2009-13 62% 2010-14 58% 2011-15 60% 59% 2012-16 2013-17 60% 2014-18 61%



Citizen Survey	/ Results	2019	2011)	1

Too High

23%(7%) Just Right

52%(56%) **Too Low**

21%(26%)

Antlerless Allocation Options							
Firearm Season Option	Increase Harvest	Stable Harvest	Decrease Harvest				
7 day antlered & 7 day concurrent	39,000	32,000	24,000				
14 day concurrent	34,000	28,000	21,000				

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	10 Characteri		
WMU	4D	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		6%	70%	22%	28%	2,743

er Harve	st												
							Ant	lered H	larvest				
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	12.000								
2005	5,591	8,354	40,000	4.7	12,000								
2006	6,776	9,878	40,000	4.0									
2007	5,765	8,073	40,000	4.9	10,000								
2008	6,593	9,310	40,000	4.2									
2009	4,971	7,192	40,000	5.6	8,000 +							_	
2010	6,321	5,472	30,052	5.6									
2011	7,144	6,561	37,000	5.7	6,000						Ш		
2012	6,922	6,325	36,000	5.7	0,000								
2013	7,165	8,225	35,000	4.3									
2014	6,461	6,832	33,000	5.0	4,000								
2015	7,240	7,197	33,000	4.6									
2016	7,921	7,234	34,000	4.7	2,000					Н	Н		
2017	10,594	8,381	34,000	4.0									
2018	8,299	8,703	34,000	3.9	_ 4				■,■,			_	
2019	8,740	10,266	46,000	4.5	pot	200 2001	2000 2009	20 2	× & .	3 4	\$ 2	6 2021	2028 2029

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	43,299
2009	62,529
2010	46,284
2011	73,017
2012	70,495
2013	67,011
2014	61,428
2015	56,905
2016	60,398
2017	63,984
2018	99,997
2019	61,822
2020	71,983




Approximately 30% of WMU is a CWD DMA in 2019

Trend

Stable

Year	Total
2008	0.34
2009	0.31
2010	0.39
2011	0.36
2012	0.37
2013	0.36
2014	0.31
2015	0.32
2016	0.31
2017	0.31
2018	0.29
2019	0.30




FOREST HEALTH

Plot - Plot Regeneration No change Plot - Plot Deer Impact No change

Regeneration Assessment Fair

Mean Deer Impact 3 or less

Year	% Adequate
2003-07	43%
2004-08	43%
2005-09	43%
2006-10	44%
2007-11	43%
2008-12	48%
2009-13	49%
2010-14	48%
2011-15	52%
2012-16	53%
2013-17	48%
2014-18	50%



Citizen Survey Results 2019 (2011)

Too High

20%(8%) Just Right

48%(46%) **Too Low**

26%(38%)

Antlerless Allocation Options								
Firearm Season Option	Increase Harvest	Stable Harvest	Decrease Harvest					
7 day antlered & 7 day concurrent	51,000	39,000	28,000					
14 day concurrent	45,000	34,000	25,000					

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	/io Characteri	Stics	
WMU	4E	% Developed	d % Forest	%Ag/Field	%Public	Area (sq mi)
		8%	54%	34%	4%	1.736

er Harve						Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹		Antiered narvest
2005	4,544	9,130	38,000	4.1	12,000	
2006	4,134	8,975	38,000	4.2		
2007	3,314	8,119	38,000	4.6	10,000	
2008	4,270	7,193	30,000	4.1		
2009	4,064	6,287	30,000	4.8	8,000	
2010	4,768	5,923	26,899	4.6		
2011	5,076	6,054	29,000	4.8	6,000	
2012	4,960	6,079	28,000	4.6	,,,,,,	
2013	6,287	7,707	26,000	3.4	4.000	
2014	5,847	5,919	21,000	3.6	4,000	
2015	6,202	6,914	25,000	3.6		
2016	7,294	7,474	25,000	3.4	2,000	
2017	8,241	8,735	27,500	3.1		
2018	6,980	9,345	32,000	3.4		┊ ┻╷┻╷┻╷┻╷┻╷┻╷┻╷┻╷┻╷┻╷┻╷┺╷
2019	7,314	9,513	34,000	3.6		or to

RED=7-day season

POST-HUNT Deer Population Trend Stable
--

Year	Total
2008	35,121
2009	37,339
2010	36,311
2011	51,706
2012	44,225
2013	48,318
2014	50,707
2015	59,206
2016	64,923
2017	62,285
2018	70,064
2019	60,055
2020	59,120




Approximately 4% of WMU is a CWD DMA in 2019

Trend

Declining

Year	Total
2008	0.44
2009	0.44
2010	0.44
2011	0.42
2012	0.45
2013	0.32
2014	0.38
2015	0.40
2016	0.35
2017	0.34
2018	0.33
2019	0.29




FOREST HEALTH

Plot - Plot Regeneration No change Plot - Plot Deer Impact No Change

Regeneration Assessment

Fair >3

Year	% Adequate
2003-07	68%
2004-08	68%
2005-09	65%
2006-10	66%
2007-11	65%
2008-12	60%
2009-13	64%
2010-14	56%
2011-15	56%
2012-16	67%
2013-17	69%
2014-18	65%



Citizen Survey Results 2019 (2011)

Too High

30%(8%) Just Right

50%(58%) **Too Low**

16%(28%)

Antlerless Allocation Options					
Firearm Season Option	Increase	Stable	Decrease		
	Harvest	Harvest	Harvest		
7 day antlered & 7 day concurrent	37,000	31,000	25,000		
14 day concurrent	36,000	30,000	24,000		

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			WIVIO Characteristics				
WMU	5A	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)	
		14%	35%	49%	11%	1.301	

er Harve	est				
					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	13,000
2005	2,396	4,690	28,000	5.8	12,000
2006	2,155	5,207	25,000	4.7	
2007	2,433	3,881	22,000	5.5	10,000
2008	2,057	3,778	19,000	4.9	
2009	2,237	4,194	19,000	4.6	8,000
2010	2,442	3,398	18,269	5.4	
2011	3,575	3,573	19,000	5.3	6,000
2012	2,795	3,596	19,000	5.3	
2013	2,825	4,098	19,000	4.6	1000
2014	2,377	3,282	19,000	5.8	4,000
2015	2,862	4,631	19,000	4.1	
2016	3,017	4,047	19,000	4.7	2,000
2017	2,925	3,811	22,000	5.7	
2018	3,091	4,649	23,000	4.9	_
2019	3,406	4,951	22,000	4.4	\$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$

RED=7-day season

POST-HUNT Deer Population	Trend	Stable

Year	Total
2008	22,602
2009	20,504
2010	20,512
2011	21,098
2012	35,598
2013	28,014
2014	29,715
2015	25,032
2016	20,081
2017	28,581
2018	33,243
2019	25,162
2020	49,801




Approximately 70% of WMU is a CWD DMA in 2019 (Former DMA1)

Tren


Stable

Year	Total
2008	0.40
2009	0.37
2010	0.43
2011	0.32
2012	0.35
2013	0.38
2014	0.34
2015	0.31
2016	0.33
2017	0.35
2018	0.28
2019	0.28



FOREST HEALTH Plot - Plot Regeneration - Plot - Plot Deer Impact - Mean Deer Impact 3 or less

Year	% Adequate
2003-07	75%
2004-08	74%
2005-09	72%
2006-10	73%
2007-11	72%
2008-12	66%
2009-13	67%
2010-14	75%
2011-15	58%
2012-16	52%
2013-17	60%
2014-18	65%



Citizen Survey Results	2019 (2011)	Too High	19%(5%)	Just Right	53%(58%) Too Low	23%(25%)

Antlerless Allocation Options					
Firearm Season Option	Increase Harvest	Stable Harvest	Decrease Harvest		
7 day antlered & 7 day concurrent	29,000	23,000	16,000		
14 day concurrent	26,000	21,000	15,000		

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.


			VVIV	io characteri	Stics	
WMU	5B	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)
		19%	28%	49%	2%	2.640

								Antle	red Ha	rvest					
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹	13.000										
2005	7,381	11,717	56,000	4.6	12,000										
2006	6,995	11,384	53,000	4.5											
2007	5,974	11,143	53,000	4.6	10,000 -										П
2008	6,762	11,184	51,000	4.4									_ 1		
2009	6,007	11,321	51,000	4.5	8,000 -							_		Н	_
2010	6,902	12,543	50,812	4.1			_	_			I _				
2011	7,174	12,943	50,000	3.9	6,000										
2012	8,503	12,519	51,000	4.1	0,000										
2013	7,443	12,847	50,000	3.9											
2014	6,908	12,368	49,000	4.0	4,000 -										
2015	8,009	11,451	50,000	4.4											
2016	8,886	12,364	50,000	4.1	2,000 -									H	_
2017	8,990	12,794	57,000	4.4											
2018	9,165	14,191	58,000	4.1						Ļ₽ ,					_
2019	10,151	14,844	67,000	4.5	2	15 50° 15	ign ring	708 Y	020 ²⁰ 25	2023	3204	20 yo	\$ 20.5°	2028	<i>50</i> 5€

RED=7-day season

POST-HUNT Deer Population	Trend	Stable
---------------------------	-------	--------

Year	Total
2008	54,020
2009	59,568
2010	53,213
2011	55,951
2012	60,723
2013	75,260
2014	63,591
2015	60,538
2016	66,282
2017	73,573
2018	85,790
2019	77,893
2020	76,623




Approximately 12% of WMU is a CWD DMA in 2019 (former DMA1)

Trend

Stable

Year	Total
2008	0.44
2009	0.42
2010	0.41
2011	0.40
2012	0.42
2013	0.41
2014	0.40
2015	0.37
2016	0.37
2017	0.41
2018	0.37
2019	0.33



FOREST HEALTH

Plot - Plot Regeneration

Plot - Plot Deer Impact


Regeneration Assessment

Fair 3 or less

Year % Adequate 2003-07 53% 2004-08 52% 2005-09 48% 2006-10 46% 2007-11 47% 2008-12 52% 2009-13 54% 2010-14 38% 2011-15 55% 2012-16 51%

2013-17

2014-18



Citizen Survey Results 2019 (2011)

49%

52%

19%(13%) Just Right 51%(58%) **Too Low** Too High 20%(21%)

Antlerless Allocation Options					
	Increase	Stable	Decrease		
Firearm Season Option	Harvest	Harvest	Harvest		
7 day antlered & 7 day concurrent	72,000	60,000	49,000		
14 day concurrent	70,000	58,000	47,000		

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

			VVIV	io characteri	stics	
WMU	5C	% Developed	% Forest	%Ag/Field	%Public	Area (sq n
		27%	37%	31%	1%	1,98

Deer Harvest Antlered Harvest Antlered Antlerless Allocation Lic/Deer¹ Year 12,000 2005 71,000 3.9 7,701 17,589 79,000 2006 7,708 16,123 4.8 10,000 2007 6,526 18,864 84,000 4.3 2008 8,729 20,238 92,000 4.4 8,000 2009 7,566 23,214 113,000 4.8 9,400 23,977 121,960 4.7 2010 2011 8,928 24,234 117,000 4.4 6,000 2012 7,825 23,648 111,000 4.5 2013 8,096 21,711 103,000 4.7 4,000 2014 8,035 22,152 95,000 4.3 2015* 7,416 13,551 70,000 5.1 2,000 2016 8,328 15,643 70,000 4.4 2017 8,846 15,644 70,000 4.4 2018 7,584 16,400 70,000 4.2 2022 2013 2014 , 2015* 100 100 , 202, 2019 7,646 14,364 70,000 4.8 * WMU Boundary Change

POST-HUNT Deer Population	Trend	Stable
---------------------------	-------	--------


Year	Total	
2008		
2009		
2010	Harvest indi	ces (i.e., antlered harvest, antlerless lic/deer), not PASAK model, used to monitor population trend
2011		
2012		
2013		
2014		

Approximately 1% of WMU is a CWD DMA in 2018

Trend

Stable

Year	Total
2008	0.44
2009	0.47
2010	0.43
2011	0.46
2012	0.49
2013	0.43
2014	0.42
2015	0.40
2016	0.44
2017	0.40
2018	0.38
2019	0.32



FOREST HEALTH

Plot - Plot Regeneration

Plot - Plot Deer Impact

Regeneration Assessment

Mean Deer Impact

Year	% Adequate
2003-07	
2004-08	
2005-09	
2006-10	
2007-11	
2008-12	
2009-13	

Forest data not considered in this developed WMU

Citizon	Survoy	Results	2010	2011\
CHUZEH	Juivev	nesuits	ZU13 I	ZUIII

Too High

33%(30%) Just Right 51%(55%) Too Low

8%(9%)

Antlerless Allocation Options						
Increase Stable Decrease						
Firearm Season Option	Harvest	Harvest	Harvest			
14 day concurrent	79,000	70,000	61,000			

 $^{^{1}}$ - The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.

 $^{^{\}rm 2}$ - Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.

			WIVIO Characteristics				
WMU	5D	% Developed	% Forest	%Ag/Field	%Public	Area (sq mi)	
		61%	18%	11%	0%	1.327	

POST-HUNT Deer Population

2013 2014

	est					Antlered Harvest
Year	Antlered	Antlerless	Allocation	Lic/Deer ¹		
2005	1,460	4,166	20,000	4.5	12,000	
2006	1,315	4,074	20,000	4.7		
2007	977	5,185	20,000	3.8	10,000	
2008	1,343	4,533	22,000	4.7		
2009	1,130	3,911	22,000	5.2	8,000 -	
2010	1,144	3,721	22,000	5.1		
2011	1,156	3,827	22,000	4.7	6,000	
2012	1,325	3,766	19,000	4.7		
2013	1,589	4,483	18,000	4.0	4,000	
2014	1,317	3,788	18,000	4.7	4,000	
2015*	2,191	5,172	24,000	4.6		_ 1 1
2016	2,908	6,452	30,000	4.6	2,000	
2017	3,327	7,526	30,000	3.9		
2018	2,631	6,001	28,000	4.6	- +	
2019	2,488	6,721	29,000	4.3	,	

Year	Total	
2008		
2009		
2010	Harvest ind	j
2011		
2012		


Trend

Stable

Trend

Stable

Total
0.41
0.49
0.48
0.49
0.44
0.41
0.48
0.42
0.42
0.40
0.46
0.33



FOREST HEALTH Plot - Plot Regeneration Plot - Plot Deer Impact Mean Deer Impact

Year	% Adequate
2003-07	
2004-08	
2005-09	
2006-10	
2007-11	
2008-12	
2009-13	

Citizen Survey Results	2019 (2011)	Too High	33%(25%) Just Right	51%(55%) Too Low	8%(18%)

Antlerless Allocation Options					
Increase Stable					
Firearm Season Option	Harvest	Harvest	Harvest		
14 day concurrent	35,000	29,000	23,000		

- 1 The number of antlerless licenses sold that it takes to harvest an antlerless deer. The number sold will differ from the allocation.
- $^{\rm 2}$ Harvest fawn to doe ratio is calculated as percent of fawns in the antlerless harvest.